



**Effect of the presence of neomycin in goat's milk on the making and characteristics of Tronchón cheese**

Quintanilla P., Beltrán M.C., Palau B., Escriche M.I., Molina M.P.

*in*

Ruiz R. (ed.), López-Francos A. (ed.), López Marco L. (ed.).  
Innovation for sustainability in sheep and goats

Zaragoza : CIHEAM

Options Méditerranéennes : Série A. Séminaires Méditerranéens; n. 123

2019

pages 283-286

Article available on line / Article disponible en ligne à l'adresse :

<http://om.ciheam.org/article.php?IDPDF=00007899>

To cite this article / Pour citer cet article

Quintanilla P., Beltrán M.C., Palau B., Escriche M.I., Molina M.P. **Effect of the presence of neomycin in goat's milk on the making and characteristics of Tronchón cheese.** In : Ruiz R. (ed.), López-Francos A. (ed.), López Marco L. (ed.). *Innovation for sustainability in sheep and goats*. Zaragoza : CIHEAM, 2019. p. 283-286 (Options Méditerranéennes : Série A. Séminaires Méditerranéens; n. 123)



CIHEAM  
Centre International de Hautes Études  
Agronomiques Méditerranéennes

<http://www.ciheam.org/>

<http://om.ciheam.org/>

# Effect of the presence of neomycin in goat's milk on the making and characteristics of Tronchón cheese

P. Quintanilla<sup>1</sup>, M.C. Beltrán<sup>1</sup>, B. Palau<sup>1</sup>, M.I. Escriche<sup>2</sup> and M.P. Molina<sup>1</sup>

<sup>1</sup>Institute for Animal Science and Technology

<sup>2</sup>Institute of Food Engineering for Development. Universitat Politècnica de València.  
Camino de Vera, s/n. 46022, Valencia, Spain

---

**Abstract.** Antibiotic residues in milk and dairy products are a potential problem for public health, and for the dairy industry. This study investigated the effect of the presence of neomycin in goat's milk on the production and characteristics of semi-hard Tronchón cheese. Cheeses were made, in duplicate, from raw goat's milk without antibiotics and with addition of neomycin at the Maximum Residue Limit (1500 µg/kg). Possible changes in the characteristics and the presence of antibiotic residues in the cheeses were evaluated at different time points along ripening (1, 30 and 60 days). Processing time was increased ( $48 \pm 18$  min) by the presence of neomycin in milk. Regarding cheeses, lower values of water activity as well as reduced free fatty acids and free amino-acid concentrations were detected in cheeses from milk with the antibiotic, suggesting a minor biochemical activity throughout ripening. Thus, a greater lightness and lower redness value were observed in this type of cheese, also presenting lower springiness. Moreover, high concentrations of neomycin were retained in the cheese and concentrated during ripening, leading to a concentration factor of 4.9, with respect to the initial concentration in the milk, potentially implying a risk to public health.

**Keywords.** Antibiotic – Neomycin – Goat milk – Cheese.

## **Effet de la présence de néomycine dans le lait de chèvre sur la fabrication et les caractéristiques du fromage Tronchón**

**Résumé.** La présence de résidus d'antibiotiques dans les produits laitiers peut constituer un problème pour la santé publique et pour l'industrie laitière. Dans ce travail, nous avons évalué l'effet de la présence de néomycine dans le lait de chèvre sur la fabrication et les caractéristiques du fromage Tronchón. Les fromages ont été fabriqués, en double exemplaire, à partir du lait de chèvre cru sans antibiotiques et avec l'addition de néomycine à la limite maximale de résidu (1500 µg/kg). Nous avons étudié les changements dans les caractéristiques du fromage et la présence de résidus à différentes périodes de maturation (1, 30 et 60 jours). La durée du procédé de fabrication augmentait ( $48 \pm 18$  min) sous l'effet de la présence de néomycine dans le lait de chèvre. Concernant les caractéristiques du fromage, nous avons observé une diminution dans les valeurs de l'activité de l'eau ainsi qu'une réduction dans les concentrations d'acides gras libres et d'acides aminés libres dans les fromages avec l'antibiotique, en suggérant une activité biochimique mineure pendant la maturation. Nous avons aussi observé une plus grande légèreté, moins rougeur et une plus faible élasticité dans ce type de fromage. En plus, pendant la maturation, les concentrations de néomycine dans le fromage augmentaient 4,9 fois par rapport à la concentration initiale dans le lait, ce qui pourrait impliquer un risque pour la santé publique.

**Mots-clés.** Antibiotique – Néomycine – Lait de chèvre – Fromage.

---

## **I – Introduction**

Goat's milk demand has increased in recent years as it is easier to digest and is an ideal substitute for cow milk in the diet of children, the elderly and people with allergies to cow milk (Haenlein, 2004). Traditionally, goat's milk production was intended for the manufacture of dairy products, especially cheese. In dairy goats, the use of antibiotics to treat mastitis and other animal health con-

ditions is a common practice that might contaminate the milk supply if appropriate measures are not taken. Among antibiotics, the aminoglycosides (gentamicin, neomycin and streptomycin, among others) are usually applied in to treat infectious diseases caused by gram-negative bacteria (Davies and Wright, 1997). The presence of antibiotic residues in milk can affect consumer health, either generating resistance to antibiotics or causing allergies (EFSA, 2016), as well as the production, thus affecting the growth of microorganisms responsible for fermentation or biochemical reactions (Novés *et al.*, 2015). For consumer protection, the European Union (EU Regulation 37/2010), established Maximum Residue Limits (MRLs) in products of animal origin such as milk, but so far, they have not been established for dairy products, such as butter, yogurt or cheese. It has been shown that the presence of antimicrobial residues in milk may result in the retention of varying amounts of these substances in cheese (Adetunji, 2011; Cabizza *et al.*, 2016). Also, the presence of antibiotics might have negative effects on cheese quality parameters. However, related studies are very scarce. Thus, the aim of this study was to analyse the effect of the presence of neomycin, on the production and the characteristics of Tronchón cheeses along ripening.

## II – Materials and methods

### 1. Cheese manufacture

Tronchón cheeses were made at the Universitat Politècnica de València (UPV) pilot plant using raw milk from the experimental flock of Murciano-Granadina goats of UPV. Cheese manufacture was performed in duplicate from goat's milk without antibiotics (control) and with neomycin (N1876, Sigma-Aldrich Química, SA, Madrid, Spain) at an equivalent concentration to the MRL (1,500 µg/kg) in a 50 l vat, following the traditional Tronchón cheese-making procedure. After moulding, the pressed rennet curd was salted and ripened for 60 days.

### 2. Cheese analysis

The physicochemical, colour and texture properties of the Tronchón cheeses were evaluated at 1, 30 and 60 days of ripening. Cheese composition (moisture, fat, protein and salt content) was determined in duplicate using a FoodScan infrared device (Foss, Foss Iberia, Barcelona). The water activity of samples (*aw*) was measured in duplicate by using a dew point hygrometer (Decagon Devices Inc., Aqualab 4TE, USA). Regarding pH, a pH-meter with a penetration electrode (Crison Instruments, S.A., Basic 20, Spain) was employed, making 6 measurements per sample. The free amino-acid (FFA) concentration, as indicator of proteolysis in cheeses, was determined by the method reported by Folkertsma and Fox (1992). The free fatty acid (FFA) concentration, as indicator of lipolysis level, was performed following the methodology described by Nuñez *et al.* (1986). The colour of the cheeses was measured using a Minolta spectrophotometer, model CM-3600D (Minolta, Japan), considering the coordinates of the CIE colour space  $L^* a^* b^*$ , which were obtained by reflectance, with illuminant D65 and the observer 10°. The texture was determined through a texture profile analysis (TPA) using a TA.XT.plus texturometer (Stable Micro Systems, UK). A double compression analysis was performed with a cylindrical probe of 45 mm diameter (P/45), with a deformation percentage of 50% and a holding time of 5 seconds. 9 measurements were made per cheese in cylinders 2 cm in diameter and 1 cm thick. The concentration of residual neomycin in cheeses was analysed in the Laboratory of Public Health of Galicia (Spain), using a High Performance Liquid Chromatography (HPLC-MS-MS) method.

### 3. Statistical analysis

Data generated by this study were subjected to a two-way analysis of variance (ANOVA), including the effect of antibiotic concentration (0 and 1500 µg/kg) and ripening time (1, 30 and 60 days)

as variation factors. The least significant differences were calculated using the LSD test (least significant difference) and a significance level of  $\alpha=0.05$ , using Statgraphics Centurion XVI (version 16.2.04; Statpoint Technologies, Inc. Warrenton, Virginia, VA, USA).

### III – Results and discussion

The cheese-making process was moderately affected by the presence of neomycin in goat's milk which inhibit the starter culture activity and increased by  $48 \pm 18$  min the time required to reach the final pH of the cheeses (pH = 5.3) during the pressing stage. Antibiotics affecting kinetic acidification in the manufacture of fermented products have been reported by other authors (Novés *et al.*, 2015; Cabizza *et al.*, 2017). Regarding the characteristics of the cheeses, Table 1 shows the parameters evaluated along with the results of the analysis of variance. As shown in Table 1, a lower water activity ( $p<0.001$ ) as well as shorter concentrations of FFA ( $p<0.05$ ) and FAA ( $p<0.01$ ) were found in the cheeses made from spiked goat's milk, suggesting that the microbial metabolism was also reduced along ripening by the presence of the residual amounts of neomycin in the curd. Thus, the lower degree of biochemical reactions in cheeses with the antibiotic could be related to the higher lightness ( $p<0.001$ ) observed in this type of cheese (Rohm and Jaros, 1997), which also presented a lower redness value ( $p<0.05$ ), especially, in the second month of ripening ( $p<0.001$ ). Regarding texture analysis, the presence of the antibiotic significantly affected the springiness of the cheeses, being lower ( $p<0.05$ ) in cheeses with neomycin. Cohesiveness also declined drastically during the first month of ripening ( $p<0.01$ ) in cheeses made from contaminated milk.

**Table 1. Average values of parameters analysed in Tronchón cheese and ANOVA F-ratio for each of the two factors: neomycin concentration (C) and ripening time (T) and their interaction (C\*T)**

Parameters	Neomycin concentration ( $\mu\text{g}/\text{kg}$ )			Ripening time (days)				ANOVA F-ratio		
	0	1,500	SE <sup>1</sup>	1	30	60	SE <sup>1</sup>	C	T	C*T
pH	5.29	5.27	0.02	5.36 <sup>b</sup>	5.26 <sup>a</sup>	5.22 <sup>a</sup>	0.02	0.47 <sup>ns</sup>	11.02 <sup>*</sup>	3.00 <sup>ns</sup>
Moisture (%)	40.27	40.86	0.36	42.81 <sup>c</sup>	40.64 <sup>b</sup>	38.24 <sup>a</sup>	0.44	1.38 <sup>ns</sup>	27.29 <sup>***</sup>	0.26 <sup>ns</sup>
Fat (%)	32.56	32.11	0.26	31.72 <sup>a</sup>	32.03 <sup>a</sup>	33.25 <sup>b</sup>	0.32	1.47 <sup>ns</sup>	6.06 <sup>*</sup>	0.22 <sup>ns</sup>
Protein (%)	22.24	21.79	0.19	20.76 <sup>a</sup>	21.85 <sup>b</sup>	23.45 <sup>c</sup>	0.23	3.02 <sup>ns</sup>	37.32 <sup>***</sup>	0.05 <sup>ns</sup>
NaCl (%)	1.85	1.87	0.05	1.64 <sup>a</sup>	1.85 <sup>b</sup>	2.09 <sup>c</sup>	0.06	0.07 <sup>ns</sup>	11.62 <sup>***</sup>	0.79 <sup>ns</sup>
Water activity	0.97 <sup>b</sup>	0.96 <sup>a</sup>	0.00	0.98 <sup>c</sup>	0.97 <sup>b</sup>	0.96 <sup>a</sup>	0.00	26.79 <sup>***</sup>	432.7 <sup>***</sup>	0.39 <sup>ns</sup>
FAA <sup>2</sup>	2.76 <sup>b</sup>	2.45 <sup>a</sup>	0.06	0.76 <sup>a</sup>	2.96 <sup>b</sup>	4.09 <sup>c</sup>	0.08	11.84 <sup>**</sup>	426.2 <sup>***</sup>	3.40 <sup>ns</sup>
FFA <sup>3</sup>	2.67 <sup>b</sup>	2.34 <sup>a</sup>	0.08	1.64 <sup>a</sup>	2.70 <sup>b</sup>	3.18 <sup>c</sup>	0.10	6.87 <sup>*</sup>	56.28 <sup>***</sup>	3.26 <sup>ns</sup>
L*	86.72 <sup>a</sup>	87.66 <sup>b</sup>	0.18	90.69 <sup>c</sup>	86.03 <sup>b</sup>	84.84 <sup>a</sup>	0.22	14.00 <sup>***</sup>	198.9 <sup>***</sup>	0.97 <sup>ns</sup>
a*	-1.24 <sup>b</sup>	-1.31 <sup>a</sup>	0.02	-0.32 <sup>c</sup>	-1.57 <sup>b</sup>	-1.93 <sup>a</sup>	0.03	5.19 <sup>*</sup>	962.3 <sup>***</sup>	8.23 <sup>***</sup>
b*	11.83	11.50	0.13	10.74 <sup>a</sup>	12.56 <sup>c</sup>	11.71 <sup>b</sup>	0.16	3.17 <sup>ns</sup>	31.57 <sup>***</sup>	1.62 <sup>ns</sup>
Hardness (N)	23.36	22.58	0.52	30.28 <sup>b</sup>	19.11 <sup>a</sup>	19.53 <sup>a</sup>	0.63	1.13 <sup>ns</sup>	99.5 <sup>***</sup>	0.85 <sup>ns</sup>
Adhesiveness (N*s)	-1.01	-1.04	0.03	-0.58 <sup>a</sup>	-1.24 <sup>b</sup>	-1.26 <sup>b</sup>	0.04	0.39 <sup>ns</sup>	84.5 <sup>***</sup>	2.16 <sup>ns</sup>
Springiness	0.63 <sup>b</sup>	0.60 <sup>a</sup>	0.01	0.82 <sup>c</sup>	0.56 <sup>b</sup>	0.46 <sup>a</sup>	0.01	5.80 <sup>*</sup>	619.1 <sup>***</sup>	2.81 <sup>ns</sup>
Cohesiveness	0.43	0.42	0.00	0.68 <sup>c</sup>	0.33 <sup>b</sup>	0.27 <sup>a</sup>	0.00	1.64 <sup>ns</sup>	2442 <sup>***</sup>	7.47 <sup>**</sup>
Chewiness (N)	7.92	7.52	0.20	17.00 <sup>c</sup>	3.64 <sup>b</sup>	2.51 <sup>a</sup>	0.25	1.99 <sup>ns</sup>	1062 <sup>***</sup>	0.02 <sup>ns</sup>

<sup>1</sup> SE: standard error.

<sup>2</sup> FAA: Free amino acids (mg leucine/100 g cheese).

<sup>3</sup> FFA: Free Fatty Acids (meq/100 g milk fat).

a, b, c: Different letters in the same row indicate significant differences ( $p<0.05$ ); \*\*\* $P<0.001$ ; \*\* $P<0.01$ ; \* $P<0.05$ ; ns: non-significant.

THPLC analysis indicated a high concentration of neomycin in the cheese at the beginning of maturation which increased along time (day 1:  $5,590 \pm 14 \mu\text{g/kg}$ ; day 30:  $6,420 \pm 707 \mu\text{g/kg}$ ; day 60:  $7,415 \pm 120 \mu\text{g/kg}$ ), due to the loss of moisture occurring during this period (Table 1). Similar amounts of residues were obtained along ripening ( $p > 0.05$ ) when antibiotic concentrations are expressed on dry basis. Thus, although neomycin is water soluble (Papich, 2016) results herein indicate that this substance is largely retained in rennet curd and shows high stability under the maturation conditions. In fact, a concentration factor of  $4.9 \pm 0.1$ , with respect to the initial concentration of neomycin in milk, was calculated in Tronchón cheese at the end of ripening. These results agree to those reported by Giraldo *et al.* (2017) when assessing the transfer of antibiotics from goat milk to whey in a laboratory scale study.

## IV – Conclusions

The presence of neomycin, at safety level, in goat's milk destined to the manufacture of ripened cheese may affect the development of biochemical processes during maturation leading to organoleptic shortcomings in the final product. Moreover, high concentrations of the antibiotic can be retained in mature cheese, this poses a potential risk for the consumer. It would be convenient to assess the retention mechanisms of this substance and other antibiotics in depth to avoid public health issues.

## Acknowledgments

This work is part of the AGL-2013-45147-R funded by the Ministry of Science and Innovation (Madrid, Spain). The authors thank the UPV (Program of Research and Development) for funding the PhD scholarship FPI-2014 attained by P. Quintanilla.

## References

- Adetunji V. O., 2011.** Effects of processing on antibiotic residues (Streptomycin, Penicillin-G and Tetracycline) in soft cheese and yoghurt processing lines. In: *Pakistan Journal of Nutrition*, 10, p. 792-795.
- Cabizza R., Rubattu N., Salis S., Pes M., Comunian R., Paba A. and Urgeghe P.P., 2017.** Transfer of oxy-tetracycline from ovine spiked milk to whey and cheese. In: *International Dairy Journal*, 70, p. 12-17.
- Davies J. and Wright G., 1997.** Bacterial resistance to aminoglycoside antibiotics. In: *Trends in microbiology*, 5(6), p. 234-240.
- EFSA, 2016.** The European Union summary report on antimicrobial resistance in zoonotic and indicator bacteria from humans, animals and food in 2014. EFSA Journal n° 4380:2016. European Food Safety Authority, Parma, Italia.
- Folkertsma B. and Fox P.F., 1992.** Use of the Cd-ninhydrin reagent to assess proteolysis in cheese during ripening. In: *Journal Dairy Research*, 59, p. 217-224.
- Giraldo J., Althaus R.L., Beltrán M.C. and Molina M.P., 2017.** Antimicrobial activity in cheese whey as an indicator of antibiotic drug transfer from goat milk. In: *International Dairy Journal*, 69, p. 40-44.
- Haenlein G.F.W., 2004.** Goat milk in human nutrition. In: *Small Ruminant Research*, 51, p. 155-163.
- Novés B., Librán C.M., Licón C.C., Molina M.P., Molina A. and Berruga M.I., 2015.** Technological failures caused by cephalaxine on set-type sheep's milk yogurt. In: *CyTA J. Food*, 13, p. 408-414.
- Núñez M., García-Aser C., Rodríguez-Martin M.A., Medina M. and Gaya P., 1986.** The effect of ripening and cooking temperatures on proteolysis and lipolysis in Manchego cheese. In: *Food Chemistry*, 21, p. 115-123.
- Papich M.G., 2016.** Saunders Handbook of Veterinary Drugs. Neomycin. Small and Large Animal. In: *Elsevier Health Sciences*, 4 e, p. 559-560.
- Rohm H. and Jaros D. 1997.** Colour of hard cheese. 2. Factors of influence and relation to compositional parameters. In: p. 259-264.