

## Nutritional consequences on the outcome of parasitic challenges on small ruminants

Athanasiadou S., Kyriazakis I., Giannenas I., Papachristou T.G.

in

Papachristou T.G. (ed.), Parissi Z.M. (ed.), Ben Salem H. (ed.), Morand-Fehr P. (ed.).  
Nutritional and foraging ecology of sheep and goats

Zaragoza : CIHEAM / FAO / NAGREF

Options Méditerranéennes : Série A. Séminaires Méditerranéens; n. 85

2009

pages 29-40

Article available on line / Article disponible en ligne à l'adresse :

<http://om.ciheam.org/article.php?IDPDF=800981>

To cite this article / Pour citer cet article

Athanasiadou S., Kyriazakis I., Giannenas I., Papachristou T.G. **Nutritional consequences on the outcome of parasitic challenges on small ruminants.** In : Papachristou T.G. (ed.), Parissi Z.M. (ed.), Ben Salem H. (ed.), Morand-Fehr P. (ed.). *Nutritional and foraging ecology of sheep and goats.* Zaragoza : CIHEAM / FAO / NAGREF, 2009. p. 29-40 (Options Méditerranéennes : Série A. Séminaires Méditerranéens; n. 85)



<http://www.ciheam.org/>  
<http://om.ciheam.org/>

# Nutritional consequences on the outcome of parasitic challenges on small ruminants

S. Athanasiadou\*, I. Kyriazakis\*\*\*, I. Giannenas\*\* and T.G. Papachristou\*\*\*

\*Animal Nutrition and Health Department, The Scottish Agricultural College, West Mains Road, Edinburgh, EH9 3JG, Scotland (United Kingdom)

\*\*Faculty of Veterinary Medicine, University of Thessaly, PO Box 199, 43100 Karditsa (Greece)

\*\*\*Forest Research Institute, National Agricultural Research Foundation, 57006 Vassilika, Thessaloniki (Greece)

---

**Abstract.** Host nutrition can affect the resident and incoming populations of pathogens and consequently ameliorate the detrimental consequences of an infectious challenge in small ruminants in four different ways: (i) affect the fitness of the parasite, through the ingestion of plant compounds, such as plant secondary metabolites; (ii) alter the conditions in the gut environment from beneficial to detrimental and even toxic for parasite survival; (iii) positively affect host resistance, i.e. the ability of the host to regulate gastrointestinal nematode establishment, development, fecundity and survival; and (iv) enhance host resilience to the parasitic infection. In this paper we address the effects of nutrition either directly on parasite fitness or through its influence on the environment of the parasite. We believe the nutritional consequences on parasitised hosts have not received appropriate attention so we focus on them in this paper. We first discuss how the excess or lack of nutrients can alter the physicochemical properties of the digestive tract making this environment more hostile for parasites transmitted through the oral route. We then concentrate on specific plants rich in secondary metabolites (PSM) and review evidence on the consequences of their consumption on resident or incoming nematode population. We conclude by discussing specific prerequisites for effectively using nutrition for parasite control.

**Keywords.** Gastrointestinal parasites – Nematodes – Plant secondary metabolites – Bioactive forages.

## **Manipulation nutritionnelle et conséquences sur le parasitisme chez les petits ruminants**

**Résumé.** La nutrition de l'hôte affecte les populations de pathogènes existantes et nouvelles ce qui allège les conséquences négatives de l'infestation des petits ruminants à travers quatre voies différentes : (i) elle affecte la forme du parasite, à travers l'ingestion de composés de la plante, tels que les métabolites secondaires ; (ii) elle altère l'environnement intestinal qui passe de l'état bénéfique à l'état nuisible voire toxique pour la survie des parasites ; (iii) elle affecte positivement la résistance de l'hôte, c'est-à-dire la capacité de l'hôte à réguler l'installation du nématode gastro-intestinal, le développement, la fécondité et la survie ; et (iv) elle augmente la résilience de l'hôte à l'infection parasitaire. Cet article traite des effets de la nutrition sur la forme du parasite ou sur le milieu environnemental du parasite. On pense que les conséquences nutritionnelles sur les hôtes infestés par les parasites n'ont pas été étudiées auparavant, ce qui nous a encouragés à traiter cet aspect dans cet article. Dans une première étape on a discuté comment l'excès ou le déficit en nutriments peut altérer les propriétés physicochimiques du tube digestif rendant cet environnement plus hostile pour les parasites transmis par voie orale. On s'est concentré ensuite sur les plantes riches en métabolites secondaires et les conséquences de leur consommation sur la population de nématodes résidente ou attendue. On a conclu cette revue avec une discussion des conditions spécifiques permettant une nutrition efficace destinée au contrôle parasitaire.

**Mots-clés.** Parasites gastrointestinaux – Nématode – Métabolites secondaires – Fourrage bioactif.

---

## **I – Introduction**

Small ruminants are constantly exposed to pathogens, such as gastrointestinal nematode parasites. They eventually acquire immunity to them and successfully regulate them, although their immunity may break down during certain stages of their lives. There is abundant evidence that

host's nutrition can contribute to regulating parasite populations in a host, either by acting independently or through enhancing immunity at times of increased susceptibility to parasites (for reviews see Coop and Kyriazakis, 2001; Houdijk and Athanasiadou, 2003; Sykes and Kyriazakis, 2007). Although not all mechanisms have been investigated in depth, various ways have been proposed to account for such effects of nutrition on the parasitised host. Firstly, nutrition may directly affect the fitness of the parasite, through the ingestion of plant compounds, such as plant secondary metabolites. Secondly, the excessive consumption or the lack of certain nutrients may alter the conditions in the gut environment from beneficial to detrimental and even toxic for parasite survival. Thirdly, certain nutrients, such as protein, can affect host resistance, i.e. the ability of the host to regulate gastrointestinal nematode establishment, development, fecundity and survival. Host resistance to gastrointestinal nematodes is mainly mediated through acquired immunity and thus, host nutrition has the potential to affect gastrointestinal nematodes by affecting the rate of acquisition and the degree of expression of immunity. Finally, the nutrition of the host can ameliorate the detrimental effects of the parasitic infection on production by improving the resilience to parasitic infection. A number of reviews have been published on the effects of nutrients, such as protein, on the resistance and the resilience of parasitised hosts, and they will not be considered further in this paper (Coop and Kyriazakis, 1999, 2001; Houdijk and Athanasiadou, 2003; Kyriazakis and Houdijk, 2006; Sykes and Kyriazakis, 2007).

In this paper, we concentrate on the effects of nutrition either directly on parasite fitness or through its influence on the environment of the parasite, as we believe such nutritional consequences on parasitised hosts have not received appropriate attention. We focus on the gastrointestinal nematodes, as they are the most pervasive challenge to health and welfare of small ruminants, although similar consequences may be experienced in other type of infections. Table 1 shows epidemiologically important gastrointestinal nematodes of ruminants and their main effects on parasitised hosts.

**Table 1. Epidemiologically important gastrointestinal nematodes of sheep, goats and cattle, their site of infection and main effect on the parasitised host**

	Site of infection	Nematodes	Consequences of parasitism
Sheep, goats	Abomasum	<i>Haemonchus contortus</i>	Anaemia
		<i>Teladorsagia circumcincta</i>	Reduced food intake
	Small intestine	<i>Nematodirus battus</i>	Dehydration
		<i>Trichostrongylus colubriformis</i>	Reduced food efficiency
Cattle	Abomasum	<i>Trichostrongylus vitrinus</i>	Reduced food efficiency
		<i>Haemonchus placei</i>	Anaemia
	Small intestine	<i>Ostertagia ostertagi</i>	Reduced food intake
		<i>Cooperia oncophora</i>	Reduced food efficiency
		<i>Cooperia punctata</i>	Reduced food efficiency

Source: Adapted from Houdijk and Athanasiadou (2003).

In the first part of this paper we discuss how the excess or lack of nutrients can alter the physicochemical properties of the digestive tract, making this environment more hostile for parasites transmitted through the oral route. Although most experimental evidence is from monogastrics, we use this evidence to extrapolate on relevant mechanisms in ruminants. In the second part we concentrate on specific plants rich in secondary metabolites (PSM) and review evidence on the consequences of their consumption for resident or incoming nematode populations. As many PSM are also known for their anti-nutritional properties, we further explore the prerequisite that should be met in order for parasitised hosts to benefit from the anthelmintic properties of PSM-rich plants. In some cases, interactions may occur between these two different influences on the parasitic infection. For example, certain PSM may create an unsuitable environment for the establishment and development of parasites. We will identify such interactions when evidence is available; however, we have chosen to separate the two issues in the present

review for the sake of clarity. We conclude by making recommendations on the potential incorporation of plants in nutritional regimes that can improve the health and welfare of small ruminants.

## II – Consequences of nutrition on the parasite environment

Nematodes identify and select precise locations to reside within their hosts. Although the exact reasons for the specific choice of habitat are still not clear, the specificity of the location is so consistent that can not be disputed (Sukhdeo and Bansemir, 1996). Consequently, we may assume that any changes in the environment of the parasites may affect the incoming or the resident parasite population in two ways. The first one is by influencing the ability of incoming parasites to identify their niche thereby inhibiting their abilities to feed and reproduce. The second one is by creating a hostile/toxic environment for the incoming or resident parasites, with detrimental consequences on their survival. For example, changes in the levels or types of sugars in the diets of parasitized rats resulted in parasites changing location in the gut and consequentially experiencing penalties in their growth and reproduction (Parshad *et al.*, 1980; Crompton *et al.*, 1983). Unfortunately this research did not explore the mechanisms that underlined these observations or the manner in which the specific carbohydrates affected the environment of the parasites.

The specific conditions that carbohydrates create in the gastrointestinal tract of parasitised hosts attracted attention recently from Petkevicius *et al.* (2004, 2007). They noticed that changes in fermentation patterns in the large intestine of pigs altered pH in various compartments of the gastrointestinal tract and reduced parasite populations. They showed that the consumption of diets high in inulin, a highly degradable fructose polymer, reduced worm numbers and egg counts in pigs infected with the large intestine nematode *Trichuris suis* (Petkevicius *et al.*, 2007). Similarly, infusion of short chain fatty acids (SCFA) and lactate acid (LA), which are the fermentation products of inulin, also reduced populations of *Oesophagostomum dentatum* in pigs (Petkevicius *et al.*, 2004). The specific mode of action of the fermentation products in regulating the parasites is not clear. It was proposed that SCFA and LA may exert some direct effects on worm physiology, especially the females. High inulin diets were detrimental to parasites during the establishment of an infection and on patent infections. In addition, following the consumption of inulin and the infusion of the SCFA and LA, pH was reduced in the caecum and the proximal colon of the pig. The close association between increased levels of SCFA and LA, low pH and expulsion of nematodes suggests that changes in pH may play a major role in worm expulsion, perhaps by creating an unsuitable environment for mature and immature parasites.

To date there is no experimental evidence on whether and how the extensive dietary fibre fermentation that occurs in the rumen may influence the conditions in the gastrointestinal tract required for successful parasite establishment and survival. As with carbohydrate fermentation in monogastrics, changes in rumen pH may affect parasite establishment or survival. For example, the ecdysis of infective larvae, otherwise known as exsheathment, is a process necessary for the successful establishment in the host. Exsheathment generally occurs in the rumen for many of the gastrointestinal nematodes of small ruminants, including *Haemonchus contortus*, *Teladorsagia* spp. and *Trichostrongylus* spp. (Hertzberg *et al.*, 2002). Changes in pH of the rumen, as well as bicarbonate concentrations, influence this process. In addition, dietary changes affect the exsheathment kinetics of gastrointestinal nematodes in ruminants (De Rosa *et al.*, 2005). Dietary changes associated with changes in rumen pH, such as excessive carbohydrate ingestion, should be further considered for their antiparasitic potential. However, this should be done in combination with their antinutritional effects, modulated via negative consequences to rumen microflora, prior to advocating them as potential means for parasite control.

In addition to carbohydrate consumption, dietary protein may influence conditions in the gastrointestinal tract thereby affecting parasite survival in ruminants. This effect of protein would be

in addition to the well-established role of protein in resistance and resilience (Coop and Kyriazakis, 2001; Houdijk *et al.*, 2001). Datta *et al.* (1998) showed that protein supplementation reduced the level of parasitism in lambs and also that high-protein diets increased production of ammonia in rumen digesta. They suggest accumulated ammonia in the rumen might directly influence the rumen microenvironment by exhibiting toxic effects on *Haemonchus contortus*. High rumen ammonia also increases the pH of the rumen, which may also affect the exsheathment of the infective larvae as discussed earlier.

In addition to protein excess, extreme protein deficiency may create an unfavorable environment for parasites. Contrary to expectations, sheep fed a food low in metabolisable protein (MP) (mean: 1.7 g MP per kg body weight) had lower parasite burdens than sheep fed a food adequate in MP (Athanasiadou *et al.*, 2001b). The very low protein content of the food may have altered the structure and/or the physiology of the gut and may have created an unsuitable and hostile environment for the survival of parasites. Low-protein diets may change the morphology of the gastrointestinal mucosa (Brown *et al.*, 1963), influence local blood flow (de Aguiar-Nascimento, 2005) and increase epithelial cell apoptosis (Bodiga *et al.*, 2005), all of which may affect parasite establishment and survival. However, as none of the above hypotheses were further investigated to elucidate the possible mechanisms, it is not possible to reject or accept them. Consequently, there may be more to the role of protein in parasite regulation than the effects on resilience and resistance, and future efforts should explore the full potential of protein nutrition in parasite control.

### III – PSM-rich plants

#### 1. Anthelmintic effects

Plant secondary metabolites are organic compounds that do not seem to have a role in the primary plant metabolism, i.e. in growth, reproduction or development of plants. More than 80,000 different PSM compounds have been described, whereas more than 100,000 compounds are believed to exist in plants resulting in a great compound diversity (Acamovic and Brooker, 2005). Although PSM are present in all plants in small and moderate amounts, only a small proportion of plant species contain PSM in high amounts and those we call PSM-rich plants (Gershenson, 2002). In many cases, PSM-rich plants have anthelmintic activity, which in most cases is attributed to the PSM. Table 2 shows some examples of PSM-rich plants and the suggested anthelmintic compounds.

Saponins, lactones, glycosides and polyphenols are PSM with reported anthelmintic activity towards animal and human gastrointestinal parasites (Guarrera, 1999; Satou *et al.*, 2002). One of the PSM most systematically investigated is the polyphenolic compounds collectively called condensed tannins (CT) or proanthocyanidins. They are located inside the vacuole of the plant cell and may be present in many parts of the plant, but are usually present in higher concentrations in the woody tissues (Bernays *et al.*, 1989). Evidence of the antiparasitic activity of CT-rich plants comes from both animal experimentation and *in vitro* studies. Early observations from grazing experiments showed that CT-rich forages such as *Lotus* spp. or *Hedysarium* spp. (Niezen *et al.*, 1994, 1995, 1998) could affect Trichostrongylid infections in sheep by reducing nematode egg excretion and worm burdens. In addition, Athanasiadou *et al.* (2000a,b, 2001a) investigated CT-parasite interactions *in vivo* using quebracho (a CT-extract from *Schinopsis* spp.) as a supplement to *T. colubriformis* infected sheep. They found tannin-rich food lowered the nematode egg excretion and parasite worm burdens by 50% compared to non-tannin supplemented animals. *In vivo* data from sheep (Butter *et al.*, 2000; Niezen *et al.*, 2002; Hordegen *et al.*, 2003; Marley *et al.*, 2003), goats (Paolini *et al.*, 2003; Min *et al.*, 2004), and deer (Hoskin *et al.*, 1999) confirmed antiparasitic effects of either CT extracts or CT-rich plants. Further support of antiparasitic activity has come from *in vitro* studies where crude or purified plant extracts were tested against larval parasitic stages. Such studies, which used extracts from leguminous species such as *Lotus pedunculatus*

(greater birdsfoot trefoil), *Hedysarum coronarium* (sulla), *Onobrychis viciifolia* (sainfoin), *L. caeneata* or *Schinopsis* spp. (Molan *et al.*, 2000a,b, 2002; Athanasiadou *et al.*, 2001a; Paolini *et al.*, 2003) or woody plants, such as *Rubus fruticosus*, *Quercus robur* and *Corylus avellana* (Paolini *et al.*, 2004), all demonstrated dose-dependent anthelmintic effect against immature stages of several nematode species including *Dictyocaulus* spp., *Ostertagia* spp. as well as *H. contortus*, *T. colubriformis* and *T. circumcincta*.

**Table 2. Examples of PSM-rich plant extracts with reported anthelmintic properties, and the potential active compounds**

PSM-rich plants	Parasite	Suggested active compound (s)	Reference
<i>Catica papaya</i>	Nematodes	Papain and lysozyme	Satrija <i>et al.</i> , 1994; Stepek <i>et al.</i> , 2006
<i>Ananas comosus</i>	Nematodes	Cysteine proteinases	Stepek <i>et al.</i> , 2006
<i>Hedera helix</i>	Cestodes	Saponic extract	Julien <i>et al.</i> , 1985
<i>Macleaya cordata</i>	Nematodes	Isoquinoline alkaloids	Satou <i>et al.</i> , 2002
<i>Mallotus philippinensis</i>	Cestodes	Glycosides	Akhtar and Ahmad, 1992
<i>Chenopodium ambrosioides</i>	Nematodes	Ascaridol	Guarrera, 1999
<i>Azadirachta indica</i>	Nematodes	Tannins	Chandrawathani <i>et al.</i> , 2006
<i>Punica granatum</i>	Helminths	Polyphenols	Guarrera, 1999
<i>Artemisia vulgaris</i>	Helminths	Tannins	Guarrera, 1999
<i>Artemisia maritima</i>	Nematodes	Santonine	Waller <i>et al.</i> , 2001
<i>Paeonia suffruticosa</i>	Nematodes	Gallotannins, condensed tannins	Mohamed <i>et al.</i> , 2000
<i>Rubus idaeus</i>	Nematodes	Polyphenols	Taylor and Murant, 1966
<i>Sesbania sesban</i>	Nematodes	Condensed tannins	Ibrahim, 1992
<i>Schinopsis</i> spp.	Nematodes	Condensed tannins	Athanasiadou <i>et al.</i> , 2000a
<i>Balanites aegyptiana</i>	Nematodes	Saponin	Ibrahim, 1992
<i>Quercus robur</i>	Nematodes	Condensed tannins	Paolini <i>et al.</i> , 2004
<i>Plagiochila stephensoniana</i>	Nematodes	Polyphenols	Lorimer <i>et al.</i> , 1996
<i>Chicorium intybus</i>	Nematodes	Condensed tannins, sesquiterpene lactones	Molan <i>et al.</i> , 2003

Source: Modified from Houdijk and Athanasiadou, 2003.

The anthelmintic properties of PSM may be attributed to a number of mechanisms. Firstly, the *in vitro* results point towards a direct anthelmintic effect through inhibition of motility, feeding, development and survival (Taylor and Murant, 1966; Athanasiadou *et al.*, 2001a; Molan *et al.*, 2003; Paolini *et al.*, 2004). For example, Molan *et al.* (2000a) showed that CT extracts from greater birdsfoot trefoil, birdsfoot trefoil (*Lotus corniculatus*), sulla and sainfoin forages reduced: (i) the rate of larval development (eggs to L3 larvae); (ii) egg hatching ability; and (iii) the motility of L3 larvae. In almost all the previous *in vitro* studies, the magnitude of the anthelmintic effect was directly related to the increasing CT concentration. In some cases the activity could not be attributed to CTs alone, as crude extracts contained other compounds as well. The restoration of the inhibitory effects following the addition of specific tannin-inhibitors supports the hypotheses that at least part of the activity was due to CTs.

These *in vitro* results were supported by *in vivo* evidence, where the experimental design enabled the investigation of the specific, directly mediated anthelmintic effect in parasitised sheep (Athanasiadou *et al.*, 2000a,b, 2001a; Tzamaloukas *et al.*, 2005a). Several suggestions have been put forward to account for the possible *in vivo* anthelmintic activity of PSM. For example, CTs may

bind to L3 larvae and inhibit establishment and/or their motility (Lorimer *et al.*, 1996; Hoskin *et al.*, 1999). Furthermore, CTs may interfere with the motility of the parasite as noted from *in vitro* studies, which may inhibit development (Molan *et al.*, 2000b) or exhibit toxic effects when in direct contact with the parasite (Butter *et al.*, 2001). Electro-microscopy has shown that condensed tannins may destroy the parasite sheath making parasites more susceptible to hosts defences (Hoste *et al.*, 2006). Although there is no direct evidence, PSM may also affect the environment in which the parasites reside, for example by affecting the pH of the rumen (Villalba *et al.*, 2006; Osoro *et al.*, 2007). This is of particular importance in light of recent evidence suggesting that certain PSM delay exsheathment of the gastrointestinal nematode *H. contortus* in goats (Brunet *et al.*, 2007), which may be mediated through changes in pH in the rumen.

In addition to direct anthelmintic effects, PSM may also indirectly mediate immunological effects. PSM-rich forages may enhance immune response to parasites, either as an outcome of the PSM having immunomodulatory effects or through the nutritional value of the plant. A limited number of studies have monitored immunological parameters following the consumption of either PSM-rich forages (Tzamaloukas *et al.*, 2006) or CTs extract (Paolini *et al.*, 2003). The consumption of PSM-rich forages increased the number of mucosal mast cells and globule leukocytes, whereas CTs extracts increased the number of mast cells on the mucosa. Although it was not clear how this effect was mediated, it may have been due to the higher crude protein content of the bioactive forages compared to the grass/clover control. Consequently, parasitised hosts experienced the beneficial effects of protein supplementation on the development of resistance in parasitised hosts (Coop and Kyriazakis, 2001; Houdijk and Athanasiadou, 2003).

Furthermore, the consumption of PSM-rich plants improves resilience of parasitised hosts, i.e. their ability to cope with the parasitic infection without penalising their performance, when compared with conventionally fed animals (Niezen *et al.*, 1998; Marley *et al.*, 2006; Athanasiadou *et al.*, 2007). Although the mechanisms that underline such effects on resilience are not known, they may be related to the nutritional superiority of many PSM-rich forages. It is thus evident that the beneficial effects observed in parasitised hosts maybe the outcome of multiple actions of PSM-rich plants that are not always easily distinguished. We believe further understanding of the mechanisms that regulate their anthelmintic activity will ensure the efficient utilisation of PSM-rich plants in parasite control strategies. This is particularly relevant, because as explained in the next section, PSM are also responsible for many detrimental effects, which may further complicate their use in parasite control schemes.

## 2. Anti-nutritional effects

The consumption of PSM-rich plants has been associated with various detrimental consequences in both invertebrate and vertebrate herbivores. The severity depends on the concentration, the type and the structure of the PSM, and the concentration of other nutrients, such as proteins and carbohydrates in the diet (Mueller-Harvey, 2006). In general, herbivores that do not regularly include PSM-rich plants in their diets, such as various monogastrics and sheep (sheep generally eat plants rich in PSM) and cattle, may get more severely penalised by the consumption of PSM-rich plants. The consumption of PSM-rich plants by ruminant herbivores can result in reduced intake, weight loss, toxicity and death (Milgate and Roberts, 1995; Waghorn and McNabb, 2003). For example, condensed tannin consumption has been associated with reduced food intake, digestibility and impaired rumen metabolism (Barry and McNabb, 1999; Min *et al.*, 2003). Some types of condensed tannins are also responsible for mucosal toxicity and consequently a reduction in nutrient absorption (Reed, 1995; Dawson *et al.*, 1999; for review: Mueller-Harvey, 2006). Saponins have also reduced food intake and impaired growth (Applebaum and Birk, 1979; Milgate and Roberts, 1995), and they have additional been associated with haemolytic action and bloat in ruminants. Excessive consumption of alkaloids, glycosides and terpenoids can cause lesions in the nervous system (Conn, 1979; Mabry and Gill, 1979).

On the other hand, herbivores that usually include PSM-rich plants in their diets have developed

various mechanisms to inactivate them and overcome possible adverse effects on their health. For example, the excretion of salivary proline-rich proteins (PRP) has been considered the major physiological adaptation to neutralise excess of dietary tannin (see review by Bernays *et al.*, 1989). Deer produce PRP that have great affinity to dietary tannins (Robbins *et al.*, 1991). As tannins form complexes with PRP that pass through the digestive tract and are excreted in the faeces of herbivores (Mole *et al.*, 1990), they do not bind to proteins. Consequently, these animals may not experience negative consequences from the PSM. Other mammals, such as goats, that usually include tannin-rich grazing in their diets have developed other adaptation mechanisms. For example tannin tolerant, resistant or degrading microbial populations enable feral goats to cope with tannin-containing forages (Acamovic and Brooker, 2005). As a consequence, goats do not seem to suffer toxic effects of tannins at levels that other animals would (Silanikove *et al.*, 1996). Sheep do not have any protective/adaptive mechanisms for consuming tannins (Robbins *et al.*, 1991), and thus are more susceptible to their anti-nutritional effects. Mammals that have developed mechanisms to cope with the anti-nutritional consequences of PSM consumption may experience the anti-parasitic effects of PSM in a different manner compared to those without such mechanisms. For example, studies performed in parasitised goats suggest that the administration of Quebracho extract, a condensed tannin extract, may reduce the fecundity of abomasal nematodes (Paolini *et al.*, 2003), which contradicts the findings from sheep (Athanasiadou *et al.*, 2001a). In the latter, consumption of Quebracho extract affected the numbers and the fecundity of intestinal, but not abomasal nematodes. Figure 1 shows the total worm population recovered in sheep infected with either an intestinal (*Trichostrongylus colubriformis*) or an abomasal (*Haemonchus contortus*) nematode species following or not the administration of Quebracho extract. The fecundity of the nematodes followed a similar trend. The differences between sheep and goats may be attributed to the different physiological adaptations of sheep and goats mentioned above. Such adaptations, developed by goats to counteract secondary metabolites present in browse plants, may modulate the bioavailability of tannins, i.e. the amount of free condensed tannins available to interact with nematodes, and thus goats may experience the anthelmintic effects of tannins in a different manner from sheep. On how different hosts experience the positive and negative effects of PSM, and whether they can use them for their own benefit, is an issue that requires further investigation, prior to incorporating PSM in parasite control schemes.



**Fig. 1.** Total worm population recovered from sheep infected with either an intestinal (*Trichostrongylus colubriformis*) or an abomasal (*Haemonchus contortus*) nematode on day 35 post infection. On day 28 sheep were drenched with 8% (hatched) of food intake commercially available condensed tannin extract (Quebracho extract). Undrenched controls are also shown (solid black). The bars represent 95% confidence intervals (from Athanasiadou *et al.*, 2001a).



In order for parasitized ruminants to benefit from the anthelmintic properties of the PSM-rich plants, the antiparasitic effects should outweigh the anti-nutritional consequences on the performance of the parasitized host. The latter will depend on the severity of the consequences to the host and to the parasite, as discussed in two recent reviews (Houdijk and Athanasiadou, 2003; Athanasiadou and Kyriazakis, 2004). It will also depend on the duration of exposure to the PSM-rich plants and the species of parasitised host as discussed above. For example, parasitized hosts might be able to tolerate short-term negative consequences (e.g. toxicity) if they can attain long-term benefits (e.g. parasite reduction) (Kyriazakis and Emmans, 1992). Consequently, short-term exposure may be one option for including PSM in parasite management schemes and others will be suggested in the next part of this paper.

### 3. Suggested solutions for potential applications

Mediterranean rangelands largely consist of plants with moderate to high content of PSM, and particularly tannins (Papachristou *et al.*, 2005). Indigenous plants, such as kermes oak (*Quercus coccifera*), oriental hornbeam (*Caprinus orientalis*) and manna ash (*Fraxinus ornus*), could potentially be incorporated in low input, extensive parasite control schemes. The prerequisite is that their anti-nutritional effects are penalising the health of the herbivores to a lesser degree than does parasitism. One way of addressing this problem may be through simultaneous use of more than one PSM-rich plant. Such combined use may provide two advantages for the parasitised host. Firstly, it may dilute the potential toxic effects of the individual PSM and thus maintain the food intake of animals at high levels, providing that the different plants contain different PSM. Previous evidence has showed that lambs offered a choice of feeds that contain different PSM such as oxalates, tannins and terpenoids consume more than lambs offered a single food that contains only one of these compounds (Villalba *et al.*, 2004). Moreover, some PSM may neutralise the activity of others rendering the combination inactive, which apparently happens with tannins and saponins in mice (Freeland *et al.*, 1985), and tannins and terpenes in sheep (Mote, 2006). However, in the process of neutralising their negative activity, these interactions between different PSM may also reduce the impact of the antiparasitic activity; this is an important issue that merits further investigation.

The second benefit the parasitised host may experience from the tandem use of more than one PSM-rich plant is that their anthelmintic action may act in a synergistic and/or complementary manner and ensure better protection from the parasites compared to the one. In a recent study, the potential complementary effects of two PSM-rich forages, sula and chicory (*Chicorium intybus*) were tested in sheep parasitised with an abomasal nematode (Tzamaloukas *et al.*, 2005b). The consecutive consumption of the two PSM-rich forages did not result in a complementary effect; the authors suggest that either such an effect was not present, or it might have been masked by the immune response of the parasitised hosts. Systematic investigation is necessary to characterise the relationships among various classes of the anthelmintic PSM. The concurrent use of more than one PSM-rich plant has the potential to offer many advantages and for this reason it should not be ignored. Acquisition of such knowledge will promote the efficient and safe use of PSM-rich plants for parasite control.

An alternative way of reducing the negative effects of PSM on the parasitised host is by manipulating the availability of certain dietary nutrients. Tannin consumption reduces food intake in sheep offered a low-protein food, but not in those offered a high-protein food (Athanasiadou *et al.*, 2001b). This effect was probably mediated through the regulation of the bioavailability of the tannins in the host; more tannins were freely available to exert their negative consequences on rumen microflora and thus penalise the performance of animals on low-protein foods. On the other hand, such increased bioavailability of tannins may result in greater penalties on the parasites. In another example, protein and/or energy supplementation increased the ability of animals to eat foods that contain PSM compounds such as terpenes, tannins, oxalates, and saponins (Provenza *et al.*, 2003; Papachristou *et al.*, 2007). This was probably mediated by facilitating detoxification processes. Thus, supplementation with macro-nutrients whilst offered PSM-rich foods, can help the

animals: (i) cope better with PSMs; (ii) influence their preferences for foods high in PSMs, and (iii) likely reduce the time animals require to adapt to PSM.

As mentioned earlier, short-term grazing on PSM-rich plants may offer another possibility of minimising the negative effects of PSM, but still experiencing the positive anthelmintic effects. This way, parasitised animals would be offered access to PSM-rich plants for short periods of time, in order to reduce their level of parasitism, and then return to conventional pastures (Athanasidou and Kyriazakis, 2004). This option would be extremely attractive if effective, as potentially adverse effects of PSM consumption on the performance of parasitised hosts would only be experienced for short-term. Indeed, if short-term PSM consumption could reduce the level of parasitism, as shown for certain PSM (Paolini *et al.*, 2003; Min *et al.*, 2004; Tzamaloukas *et al.*, 2005a), then short-term penalties on animal performance could actually lead to long-term benefits. These long-term benefits would be translated as having an impact on parasite epidemiology through lower parasite excretion, exposure and greater protection from parasites in subsequent grazing seasons.

## IV – Conclusions

Host nutrition has a role in parasite management strategies as a complementary intervention to more traditional control measures, such as the use of anthelmintics. Due to the increasing rate of development of nematode resistance towards anthelmintic drugs (Jackson and Coop, 2000), as well as increasing public concerns about potential drug residues in animal products and the environment, nutrition will be required to play a more strategic role in the control of parasitism in ruminants. In the present review we have shown that through direct effects on parasite fitness, host nutrition has the potential to impact animal health, affect the epidemiology of the parasitic infection and consequently ease public concerns over environmental contamination. A strategic combination of the various consequences of host nutrition in parasitised herbivores could greatly contribute towards non-chemical, sustainable parasite control in low input, extensive systems of herbivore production. It is anticipated that nutritional control of gastrointestinal nematodes will be part of an integrated approach for parasite control, including genetic selection for increased resistance, vaccination and biological control, rather than being applied in isolation (Coop and Kyriazakis, 2001; Houdijk and Athanasidou, 2003). Thus, developments of non-chemical parasite control strategies would require a multidisciplinary approach and close collaboration among a range of specialists, including nutritionists, parasitologists, geneticists and veterinarians.

## References

- Acamovic T. and Brooker J.D., 2005.** Biochemistry of plant secondary metabolites and their effects in animals. In: *Proc. Nut. Soc.*, 64. p. 403-412.
- Akhtar M.S. and Ahmad I., 1992.** Comparative efficacy of *Mallotus philippinensis* fruit (Kamala) or Nilzan drug against gastrointestinal cestodes in Beetal goats. In: *Small Ruminant Res.*, 8. p. 121-128.
- Applebaum S.W. and Birk Y., 1979.** Saponins. In: Rosental G.A. and Janzen T.H. (eds). *Herbivores: Their Interactions with Secondary Plant Metabolites*. London: Academic Press. p. 539-566.
- Athanasidou S., Gray D., Younie D., Tzamaloukas O., Jackson F. and Kyriazakis I., 2007.** The ewes of chicory for parasite control in organic ewes and their lambs. In: *Parasitol.*, 134. p. 299-307.
- Athanasidou S. and Kyriazakis I., 2004.** Plant secondary metabolites: Antiparasitic effects and their role in ruminant production systems. In: *Proc. Nut. Soc.*, 63. p. 631-639.
- Athanasidou S., Kyriazakis I., Jackson F. and Coop R.L., 2000a.** Consequences of long-term feeding with condensed tannins on sheep parasitised with *Trichostrongylus colubriformis*. In: *Int. J. Parasitol.*, 30, 1025-1033.
- Athanasidou S., Kyriazakis I., Jackson F. and Coop R.L., 2000b.** Effects of short-term exposure to condensed tannins on adults *Trichostrongylus colubriformis*. In: *Vet. Rec.*, 146. p. 728-732.
- Athanasidou S., Kyriazakis I., Jackson F. and Coop R.L., 2001a.** Direct anthelmintic effects of condensed tannins towards different gastrointestinal nematodes of sheep: *In vitro* and *in vivo* studies. In: *Vet. Parasitol.*, 99. p. 205-219.
- Athanasidou S., Kyriazakis I., Jackson F. and Coop R.L., 2001b.** The effects of condensed tannins supplementation of foods with different protein content on parasitism, food intake and performance of sheep infected with *Trichostrongylus colubriformis*. In: *Brit. J. Nut.*, 86. p. 697-706.

- Barry T.N. and McNabb W.C., 1999.** The implications of condensed tannins on the nutritive value of temperate forages fed to ruminants. In: *Brit. J. Nutr.*, 81. p. 263-272.
- Bernays E.A., Driver G.C. and Bilgener M., 1989.** Herbivores and plant tannins. In: *Adv. Ecol. Res.*, 19. p. 263-302.
- Bodiga V.L., Boindala S., Putcha U., Subramaniam K. and Manchala R., 2005.** Chronic low intake of proteins or vitamins increases the intestinal epithelial cell apoptosis in Wistar/NIN rats. In: *Nutrition*, 21. p. 949-960.
- Brown H.O., LeVive M.L. and Lipkin M., 1963.** Inhibition of intestinal cell renewal and migration induced by starvation. In: *American J. Physiol.*, 205. p. 868-872.
- Brunet S., Aufrere J., El Babil F., Fouraste I. and Hoste H., 2007.** The kinetics of exsheathment of infective nematode larvae is disturbed in the presence of a tannin-rich plant extract (sainfoin) both *in vitro* and *in vivo*. In: *Parasitology*, 134(9). p. 1253-1262.
- Butter N.L., Dawson J.M., Wakelin D. and Buttery P.J., 2000.** Effect of dietary tannin and protein concentration on nematode infection (*Trichostrongylus colubriformis*) in lambs. In: *J. Agr. Sci.*, 134. p. 89-99.
- Butter N.L., Dawson J.M., Wakelin D. and Buttery P.J., 2001.** Effect of dietary condensed tannins on gastrointestinal nematodes. In: *J. Agr. Sci.*, 137. p. 461-469.
- Chandrawathani P., Chang K.W., Nurulaini R., Waller P.J., Adnan M., Zaini C.M., Jamnah O., Khadijah S. and Vincent N., 2006.** Daily feeding of fresh neem leaves (*Azadirachta indica*) for worm control in sheep. In: *Trop. Biomed.*, 23. p. 23-30.
- Conn E.E., 1979.** Cyanide and cyanogenic glycosides. In: Rosental G.A. and Janzen T.H. (eds). *Herbivores. Their Interactions with Secondary Plant Metabolites*. London: Academic Press. p. 387-412.
- Coop R.L. and Kyriazakis I., 1999.** Nutrition-parasite interaction. In: *Vet. Parasitol.*, 84. p. 187-204.
- Coop R.L. and Kyriazakis I., 2001.** Influence of host nutrition on the development and consequences of nematode parasitism in ruminants. In: *Trends Parasitol.*, 17. p. 325-330.
- Crompton D.W., Keymer A., Singhvi A. and Nesheim M.C., 1983.** Rat dietary fructose and the intestinal distribution and growth of *Moniliformis* (Acanthocephala). In: *Parasitology*, 86. p. 57-71.
- Datta F., Nolan J., Rowe J. and Gray G., 1998.** Protein supplementation improves the performance of parasitized sheep fed a straw-based diet. In: *Int. J. Parasitol.*, 28. p. 1269-1278.
- Dawson J.M., Buttery P.J., Jenkins D., Wood C.D. and Gill M., 1999.** Effects of dietary Quebracho tannin on nutrient utilisation and tissue metabolism in sheep and rats. In: *J. Sci. Food Agr.*, 79. p. 1423-1430.
- De Aguiar-Nascimento J.E., 2005.** The role of macronutrients in gastrointestinal blood flow. In: *Curr. Opin. Clin. Nutr. Metab. Care*, 8. p. 552-556.
- De Rosa A.A., Chirgwin S.R., Fletcher J., Williams J.C. and Klei T.R., 2005.** Exsheathment of *Ostertagia ostertagi* infective larvae following exposure to bovine rumen contents derived from low and high roughage diets. In: *Vet. Parasitol.*, 20. p. 77-81.
- Freeland W.J., Calcott P.H. and Geiss D.P., 1985.** Allelochemicals, minerals and herbivore population size. In: *Biochem. Syst. Ecol.*, 13. p. 195-206.
- Gershenson J., 2002.** Secondary metabolites and plant defences. In: Taiz L. and Zeiger E. (eds). *Plant Physiology*. Sunderland, MA, USA: Sinauer Associates. p. 283-308.
- Guarrera P.M., 1999.** Traditional anthelmintic, antiparasitic and repellent uses of plants in Central Italy. In: *J. Ethnopharmacol.*, 68. p. 183-192.
- Hertzberg H., Huwyler U., Kohler L., Rehbein S. and Wanner M., 2002.** Kinetics of exsheathment of infective ovine and bovine strongylid larvae *in vivo* and *in vitro*. In: *Parasitology*, 125. p. 65-70.
- Hordegen P., Hertzberg H., Heilmann J., Langhans W. and Maurer V., 2003.** The anthelmintic efficacy of five plant products against gastrointestinal trichostrongylids in artificially infected lambs. In: *Vet. Parasitol.*, 117. p. 51-60.
- Hoskin S.O., Barry T.N., Wilson P.R., Charleston W.A.G. and Hodgson J., 1999.** Effects of reducing anthelmintic input upon growth and faecal egg and larval counts in young farmed deer grazing chicory (*Chicorium intybus*) and perennial ryegrass (*Lolium perenne*)/white clover (*Trifolium repens*) pasture. In: *J. Agr. Sci.*, 132. p. 335-345.
- Hoste H., Jackson F., Athanasiadou S., Thamsborg S.M. and Hoskin S.O., 2006.** The effects of tannin-rich plants on parasitic nematodes in ruminants. In: *Trends Parasitol.*, 22. p. 253-261.
- Houdijk J.G.M. and Athanasiadou S., 2003.** Direct and indirect effects of host nutrition on ruminant gastrointestinal nematodes. In: Mannelje L.t., Ramirez-Aviles L., Sandoval-Castro C.A. and Ku-Vera J.C. (eds). *Matching Herbivore Nutrition to Ecosystems Biodiversity*. Mérida, Yucatán, México: Universidad Autónoma de Yucatán. p. 213-236.
- Houdijk J.G.M., Jessop N.S. and Kyriazakis I., 2001.** Nutrient partitioning between reproductive and immune functions in animals. In: *Proc. Nutr. Soc.*, 60. p. 515-525.
- Ibrahim A.M., 1992.** Anthelmintic activity of some Sudanese anthelmintic plants. In: *Phytother. Res.*, 6. p. 155-157.

- Jackson F. and Coop R.L., 2000.** The development of anthelmintic resistance in sheep nematodes. In: *Parasitology*, 120. p. S95-S107.
- Julien J., Gasquet M., Maillard C., Balansard G. and Timon-David P., 1985.** Extracts of the ivy plant, *Hedera helix*, and their anthelmintic activity on liver flukes. In: *Plant Medica*, 3. p. 205-208.
- Kyriazakis I. and Emmans G.C., 1992.** Selection of a diet by growing pigs given choices between foods differing in contents of protein and rapeseed meal. In: *Appetite*, 19. p. 121-132.
- Kyriazakis I. and Houdijk J.G.M., 2006.** Immunonutrition: Nutritional control of parasites. In: *Small Ruminant Res.*, 62. p. 79-82.
- Lorimer S.D., Perry N.B., Foster L.M. and Burgess E.J., 1996.** A nematode larval motility inhibition assay for screening plant extracts and natural products. In: *J. Agr. Food Chem.*, 44. p. 2842-2845.
- Mabry T.J. and Gill J.E., 1979.** Sesquiterpenes lactones and other terpenoids. In: Rosental G.A. and Janzen T.H. (eds). *Herbivores. Their Interactions with Secondary Plant Metabolites*. London: Academic Press. p. 501-537.
- Marley C.L., Cook R., Barrett J. Keatinge R. and Lampkin N.H., 2006.** The effects of birdsfoot trefoil (*Lotus corniculatus*) and chicory (*Cichorium intybus*) when compared with perennial ryegrass (*Lolium perenne*) on ovine gastrointestinal parasite development, survival and migration. In: *Vet Parasitol.*, 138. p. 280-90.
- Marley C.L., Cook R., Keatinge R., Barrett J. and Lampkin N.H., 2003.** The effect of birdsfoot trefoil (*Lotus corniculatus*) and chicory (*Cichorium intybus*) on parasite intensities and performance of lambs naturally infected with helminth parasites. In: *Vet. Parasitol.*, 112. p. 147-155.
- Milgate J. and Roberts D.C.K., 1995.** The nutritional and biological significance of saponins. In: *Nutr. Res.*, 15. p. 1223-1249.
- Min B.R., Barry T.N., Attwood G.T. and McNabb W.C., 2003.** The effect of condensed tannins on the nutrition and health of ruminants fed fresh temperate forages: A review. In: *Anim. Feed Sci. Tech.*, 106. p. 3-19.
- Min B.R., Pomroy W.E., Hart S.P. and Sahlou T., 2004.** The effect of short-term consumption of a forage containing condensed tannins on gastro-intestinal nematode parasite infections in grazing wether goats. In: *Small Ruminant Res.*, 51. p. 279-83.
- Mohamed A.S.A., Mori T., Islam S.Q., Sato M. and Yamasaki T., 2000.** Lethal activity of gallo- and condensed tannins against the free-living soil-inhabiting nematode *Caenorhabditis elegans*. In: *J. Pestic. Sci.*, 25. p. 410-415.
- Molan A.L., Duncan A., Barry T.N. and McNabb W.C., 2003.** Effects of condensed tannins and sesquiterpene lactones extracted from chicory on the viability of deer lungworm larvae. In: *Proc. New Zeal. Soc. Anim. Prod.*, 60. p. 26-29.
- Molan A.L., Hoskin S.O., Barry T.N. and McNabb W.C., 2000a.** Effect of condensed tannins extracted from four forages on the viability of the larvae of deer lungworms and gastrointestinal nematodes. In: *Vet. Rec.*, 147. p. 44-48.
- Molan A.L., Waghorn G.C. and McNabb W.C., 2002.** Effect of condensed tannins on egg hatching and larval development of *Trichostrongylus colubriformis* *in vitro*. In: *Vet Rec.*, 150. p. 65-69.
- Molan A.L., Waghorn G.C., Min B.R. and McNabb W.C., 2000b.** The effects of condensed tannins from seven herbage on *Trichostrongylus colubriformis* larval migration *in vitro*. In: *Folia Parasitol.*, 47. p. 39-44.
- Mole S., Rogler J.C., Morell C.J. and Butler L.G., 1990.** Herbivore growth reduction by tannins: Use of Waldbauer ratio technique and manipulation of salivary protein production to elucidate mechanisms of action. In: *Biochem. Syst. Ecol.*, 18. p. 183-197.
- Mote T.E., 2006.** Importance of foraging context and feeding sequence on preference for toxin-containing foods by lambs. MSc Thesis: Utah State University (Logan, Utah, USA).
- Mueller-Harvey I., 2006.** Unravelling the conundrum of tannins in animal nutrition and health. In: *J. Sci. Food Agr.*, 86. p. 201-2037.
- Niezen J.H., Charleston W.A.G., Robertson H.A., Shelton D. Waghorn G.C. and Green R., 2002.** The effect of feeding sulla (*Hedysarum coronarium*) or lucerne (*Medicago sativa*) on lamb parasite burdens and development of immunity to gastrointestinal nematodes. In: *Vet. Parasitol.*, 105. p. 229-245.
- Niezen J.H., Waghorn T.S., Charleston W.A.G. and Waghorn G.C., 1995.** Growth and gastrointestinal nematode parasitism in lambs grazing either lucerne (*Medicago sativa*) or sulla (*Hedysarum coronarium*) which contains condensed tannins. In: *J. Agr. Sci.*, 125. p. 281-289.
- Niezen J.H., Waghorn G.C. and Charleston W.A.G., 1998.** Establishment and fecundity of *Ostertagia circumcincta* and *Trichostrongylus colubriformis* in lambs fed lotus (*Lotus pedunculatus*) or perennial ryegrass (*Lolium perenne*). In: *Vet. Parasitol.*, 78. p. 13-21.
- Niezen J.H., Waghorn T.S., Raufautk K., Robertson H.A. and McFarlane R.G., 1994.** Lamb weight gain and faecal egg count when grazing one of seven herbage and dosed with larvae for six weeks. In: *Proc. New Zealand Soc. Anim. Prod.*, 54. p. 15-18.
- Osoro K., Mateos-Sanz A., Frutos P., García U., Ortega-Mora L.M., Ferreira L.M.M., Celaya R. and Ferre I., 2007.** Anthelmintic and nutritional effects of heather supplementation on Cashmere goats grazing perennial ryegrass-white clover pastures. In: *J. Anim. Sci.*, 85. p. 861-870.

- Paolini V., Bergeaud J.P., Grisez C., Prevot F., Dorchies P. and Hoste H., 2003.** Effects of condensed tannins on goats experimentally infected with *Haemonchus contortus*. In: *Vet. Parasitol.*, 113. p. 253-261.
- Paolini V., Fouraste I. and Hoste H., 2004.** *In vitro* effects of three woody plant and sainfoin extracts on 3rd-stage larvae and adult worms of three gastrointestinal nematodes. In: *Parasitology*, 129. p. 69-77.
- Papachristou T.G., Dziba L.E. and Provenza F.D., 2005.** Foraging ecology of goats and sheep on wooded rangelands. In: *Small Ruminant Res.*, 59. p. 141-156.
- Papachristou T.G., Dziba L.E., Villalba J.J. and Provenza F.D., 2007.** Patterns of diet mixing by sheep offered foods varying in nutrients and plant secondary compounds. In: *Appl. Anim. Behav. Sci.*, 108. p. 68-80.
- Parshad V.R., Crompton D.W.T. and Nesheim M.C., 1980.** The growth of moniliformis (Acanthocephala) in rats fed on various monosaccharides and disaccharides. In: *Proc. Royal Soc. Brit.*, 209. p. 299-315.
- Petkevicius S., Murrell K.D., Bach Knudsen K.E., Jorgensen H., Roepstorff A., Laue A. and Wachmann H., 2004.** Effects of short chain fatty acids and lactic acids on survival of *Oesophagostomum dentatum* in pigs. In: *Vet. Parasitol.*, 122. p. 293-301.
- Petkevicius S., Thomsen L.E., Bach Knudsen K.E., Murrell K.D. and Roepstorff A., 2007.** The effect of inulin on new and patent infections of *Trichuris suis* in growing pigs. In: *Parasitology*, 134. p. 121-127.
- Provenza F.D., Villalba J.J., Dziba L.E., Atwood S.B. and Banner R.E., 2003.** Linking herbivore experience, varied diets, and plant biochemical diversity. In: *Small Ruminant Res.*, 49. p. 257-274.
- Reed J.D., 1995.** Nutritional toxicology of tannins and related polyphenols in forage legumes. In: *J. Anim. Sci.*, 73. p. 1516-1528.
- Robbins C.T., Hagerman A.E., Austin P.J., McArthur C. and Hanley T.A., 1991.** Variation in mammalian physiological responses to a condensed tannin and its ecological implications. In: *J. Mamm.*, 72. p. 480-486.
- Satou T., Akao N., Matsuhashi R., Koike K., Fujita K. and Nikaido T., 2002.** Inhibitory effect of isoquinoline alkaloids on movement of second-stage larvae of *Toxocara canis*. In: *Biol. Pharmac. Bull.*, 25. p. 1651-1654.
- Satrija F., Nansen P., Bjorn H., Murtini S. and He S., 1994.** Effects of papaya latex against *Ascaris suum* naturally infected pigs. In: *J. Helminthol.*, 68. p. 343-346.
- Silanikove N., Gilboa N., Perevolotsky A. and Nitsan Z., 1996.** Goats fed tannin containing leaves do not exhibit toxic syndromes. In: *Small Ruminant Res.*, 21. p. 195-201.
- Steppek G., Lowe A.E., Buttle D.J., Duce I.R. and Behnke J.M., 2006.** *In vitro* and *in vivo* anthelmintic efficacy of plant cysteine proteinases against the rodent gastrointestinal nematode, *Trichuris muris*. In: *Parasitology*, 132. p. 681-689.
- Sukhdeo M.V.K. and Bansemir A.D., 1996.** Critical resources that influence habitat selection decisions by gastrointestinal helminth parasites. In: *Int. J. Parasitol.*, 26. p. 483-498.
- Sykes A.R. and Kyriazakis I., 2007.** Opportunities to control herbivore nematodes through manipulation of the grazing environment. In: Meng Q.X., Ren L.P. and Cao Z.J. (eds). *Proc. 7th International Symposium on the Nutrition of Herbivores: Herbivore Nutrition for the Development of Efficient, Safe and Sustainable Livestock Production*, Beijing (China), 2007. Beijing, China: China Agricultural University Press. p. 329-353.
- Taylor C.E. and Murant A.F., 1966.** Nematicidal activity of aqueous extracts from raspberry canes and roots. In: *Nematologica*, 12. p. 488-494.
- Tzamaloukas O., Athanasiadou S., Kyriazakis I., Jackson F. and Coop R.L., 2005a.** The consequences of short-term grazing of bioactive forages on established adult and incoming larvae populations of *Teladorsagia circumcincta* in lambs. In: *Int. J. Parasitol.*, 35. p. 329-335.
- Tzamaloukas O., Athanasiadou S., Kyriazakis I., Jackson F. and Coop R.L., 2005b.** Effects of sequential short-term grazing on different bioactive forages on viability and fecundity of established adult population of *Teladorsagia circumcincta*. In: *Proc. Annual Conference of the Brit. Soc. Anim. Sci.*, York (UK), 4-6 April 2005. p. 140.
- Tzamaloukas O., Athanasiadou S., Kyriazakis I., Huntley J.F. and Jackson F., 2006.** The effect of chicory (*Cichorium intybus*) and sulla (*Hedysarum coronarium*) on larval development and mucosal cell responses of growing lambs challenged with *Teladorsagia circumcincta*. In: *Parasitology*, 132. p. 419-426.
- Villalba J.J., Provenza F.D. and Han G.-D., 2004.** Experience influences diet mixing by herbivores: Implications for plant biochemical diversity. In: *Oikos*, 107. p. 100-109.
- Villalba J.J., Provenza F.D. and Olson K.C., 2006.** Terpenes and carbohydrates source influence rumen fermentation, digestibility, intake and preference in sheep. In: *J. Anim. Sci.*, 84. p. 2463-2473.
- Waghorn G.C. and McNabb W.C., 2003.** Consequences of plant phenolic compounds for productivity and health of ruminants. In: *Proc. Nutr. Soc.*, 62. p. 383-392.
- Waller P.J., Bernes G., Thamsborg S.M., Sukura A., Richter S.H., Ingebrigtsen K. and Høglund J., 2001.** Plants as de-worming agents of livestock in the Nordic Countries: Historical perspective, popular beliefs and prospects for the future. In: *Acta Vet. Scand.*, 42. p. 31-44.