

Possible alternatives to the use of antibiotics as growth promoters. New additives

Piva G., Rossi F.

in

Brufau J. (ed.), Tacon A. (ed.).
Feed manufacturing in the Mediterranean region: Recent advances in research and technology

Zaragoza : CIHEAM
Cahiers Options Méditerranéennes; n. 37

1999
pages 83-106

Article available on line / Article disponible en ligne à l'adresse :

<http://om.ciheam.org/article.php?IDPDF=99600009>

To cite this article / Pour citer cet article

Piva G., Rossi F. **Possible alternatives to the use of antibiotics as growth promoters. New additives.** In : Brufau J. (ed.), Tacon A. (ed.). *Feed manufacturing in the Mediterranean region: Recent advances in research and technology.* Zaragoza : CIHEAM, 1999. p. 83-106 (Cahiers Options Méditerranéennes; n. 37)



<http://www.ciheam.org/>
<http://om.ciheam.org/>

Possible alternatives to the use of antibiotics as growth promoters. New additives

G. Piva and F. Rossi

Istituto di Scienze degli Alimenti e della Nutrizione (ISAN), Facoltà di Agraria,
Università Cattolica del Sacro Cuore, Via Emilia Parmense 84, 29100 Piacenza, Italy

SUMMARY - The non-therapeutic use of antibiotics may be reduced by administering both microbial cultures and molecules such as oligosaccharides and lectins. In the former case, an attempt is made at preventing intestinal pathogens from settling down, by administering microorganisms which can colonize the digestive tract and leave out all dangerous bacteria. The microorganisms mostly used in connection with monogastric animals are bacteria from the *Bacillus*, *Bifidobacterium* and *Lactobacillus* genera and yeasts from *Saccharomyces cerevisiae* genus. The use of *S. cerevisiae* and *Aspergillus oryzae* is popular among adult ruminants, leading to weight gain both in calves and bullocks. In pre-ruminant cattle, the use of lactic bacteria (*Lactobacillus* and *Bifidobacterium*) also deserves some interest. In the field of chemical probiosis, the use of fructo- and gluco-oligosaccharides, capable of selectively stimulating the lactic bacteria, has yielded very interesting results both in monogastrics and in calves. Equally promising is the dietary supplementation with oligomannans and lectins. Such compounds saturate and bind to the enterocyte receptors which are present on the cell walls of pathogenic bacteria, thus preventing them from colonizing the intestinal lumen.

Key words: Antibiotics, probiotics, additives.

RESUME - "Alternatives possibles à l'utilisation d'antibiotiques comme promoteurs de croissance. Nouveaux additifs". L'utilisation non thérapeutique des antibiotiques peut être réduite soit par des molécules comme les lectines et les oligosaccharides soit par l'administration de cultures microbiennes. Dans le premier cas on cherche à empêcher l'installation des pathogènes intestinaux en administrant des micro-organismes qui colonisent l'appareil digestif en excluant les germes dangereux. Les micro-organismes les plus utilisés chez les monogastriques appartiennent aux genres *Bacillus*, *Bifidobacterium* et *Lactobacillus* parmi les bactéries et aux *Saccharomyces cerevisiae* parmi les levures. Parmi les ruminants adultes est diffusé l'emploi de *S. cerevisiae* et de *Aspergillus oryzae*, qui ont permis les augmentations de poids soit chez les bouvillons soit chez les veaux. Chez le bovin pré-ruminant il faut mentionner l'emploi des bactéries lactiques (*Lactobacillus* et *Bifidobacterium*). Dans le domaine de la probiotique chimique l'utilisation de fructo et gluco oligosaccharides qui peuvent stimuler sélectivement les bactéries lactiques, a permis d'obtenir des résultats très intéressants chez les monogastriques et dans les veaux. L'intégration alimentaire avec des oligomannanes et lectines est également prometteuse. Les oligomannanes et les lectines sont des composés qui se lient, en les saturant, aux récepteurs pour les entérocytes présents sur les parois cellulaires des bactéries pathogènes en les empêchant de coloniser la lumière intestinale.

Mots-clés : Antibiotiques, probiotiques, additifs.

Introduction

Adequate control of the activity of microorganisms living in symbiosis with the higher animal in the digestive tract of ruminants and/or monogastrics is essential to ensure balanced biological activity, ability to react to stress under intensive farming conditions, good immune response, adequate health status, excellent reproductive performance and reduced environmental impact.

Since the early 50's, thanks to the use of antibiotics as additives promoting performance in animal nutrition, considerable improvements have been obtained in production.

The non-therapeutic use of antibiotics has the following aims:

Prophylactic purposes: (i) in young animals, to reduce the enteric diseases whose onset is favoured by incomplete development of the immune system during the first weeks of life; and (ii) in adult animals, to prevent the onset of feed-induced pathologies (acidosis in steers).

Animal breeding purposes: The lower degree of microbial contamination in the gut causes less development in the intestinal lumen (Gaskins, 1996), with reduced energy and protein requirements for re-synthesizing the enteric cells and greater availability of nutrients to be used for production purposes. In oloxic animals the enterocyte turnover is accelerated by approximately twice versus the axenic animals (Vanbelle *et al.*, 1990).

Antimicrobials have also proved to be important for sustainable livestock production and for the control of animal infections that could be passed on to humans. However, concerns about using antibiotics in livestock is growing. In 1995, the WHO set up a special working group to assess this problem. More recently, the treatment of enteric diseases in children living in developing countries was found to be increasingly difficult, due to the spreading of resistance to antibiotics among the human intestinal pathogens.

"The magnitude of the medical and public health impact of antimicrobial use in food animal production is not known. It is unrefuted that the use of antimicrobials leads to the selection of resistant bacteria and that the scope of the emerging problem depends, among other things, on duration of exposure to and concentration of the antimicrobial. Residues of antimicrobial agents in food of animal origin in excess of the agreed acceptable minimum residue levels (MRLs) may contribute to generation of resistance in bacteria in humans. However the current evidence suggests that the risk is low. Of more concern may be that such residues could indicate inappropriate use of antimicrobials by producer. The medical consequences of resistance acquisition in bacteria of animal origin are highlighted by the following examples.

Salmonella-Campylobacter - Following the introduction of fluoroquinolones for use in food-producing animals the emergence of *Salmonella* serotypes and *Campylobacter jejuni* with reduced susceptibility to fluoroquinolones in humans has become a cause of particular concern.

Enterococci - The increase of glycopeptide-resistant enterococci from animals, linked to the use of some glycopeptides growth promoter (e.g., avoparcine) can reach humans via the food chain. Glycopeptide-resistant enterococci cause serious infections in hospitalized immune-impaired patients. There is concern that there will be increased dissemination of glycopeptide resistance genes to *Enterococcus faecalis* and their spread to other gram-positive organisms, particularly to multiresistant *Staphylococcus aureus* for which vancomycin is the drug last resort. This medical impact would be greatest in countries where vancomycin is used intensively; in the United States of America for instance vancomycin is used intensively and avoparcine has never been used.

Escherichia coli - Multiresistant *E. coli* have been selected by the use of broad spectrum antimicrobials in both livestock and humans. The development of antimicrobial resistance in *E. coli* creates problems due to their high propensity to disseminate antimicrobial resistance genes. Resistance genes have been traced from *E. coli* in animals to *E. coli* in humans. Certain *E. coli* are foodborne pathogens.

Because of the growing global need for food and potential public health consequences of the transmission of resistant bacteria through the food chain, the objectives for risk management at the animal production level are to assure the efficient production of safe and wholesome food of animal origin for human consumption and to reduce potential public health risks associated with farming practices to enable the growth of the global food supply" (from WHO meeting Berlin, Germany, 13-17 October 1997). Recently, (February 2, 1988) the Swedish Government requested a ban for the non therapeutic use of antibiotics.

However, one cannot possibly think of imposing a sudden and generalized ban on such products since this would imply severe repercussions on the food supplies to the poorest countries. The FAO has in fact estimated that, should antibiotics be banned from animal breeding as performance promoters, one would be faced with a 30% reduction in the availability of proteins from animal origin throughout the World, which would certainly make the state of chronic malnutrition in developing countries even worse.

With the appearance of the first problems related to the use of antibiotics, the possibility of using probiotics as growth promoters or with prophylactic effects began to be seriously considered for

scientific purposes. Probiotics are "*live microorganisms capable of inducing a beneficial effect on the balance of microorganisms in the digestive tract*".

With Directive 93/113, the EU acknowledged the validity of this category of products as "additives", in that they improve the production performance and the quality of products of animal origin.

One of the most important advantages of such additives is the absence of undesired antibiotics residues in meat, milk and eggs.

A first tentative list of products authorized in the different EU countries is reported in the Official Journal of the European Community of September 11, 1996 - C 263. Apart from positive results, the use of microorganisms has at times yielded a disappointing response. The method of administration has in fact very much affected the results. A critical point was, above all, the possibility of using a substrate (feedstuff) capable of favouring the development of useful microorganisms and affecting the adhesion of undesired or pathogenic microorganisms to the gut walls. It is not only a matter of mechanically controlling the competition between microorganisms, shifting it towards those that are regarded as useful. It is rather a question of acting on the metabolism of microorganisms and of the gut wall, in order to control the metabolic activity of the digestive tract as a whole. It is therefore necessary to act nutritionally on the digestive tract microorganisms by way of increasing the useful microorganisms, perhaps via their exogenous supply, at the same time providing the compounds which may favour competition with the undesired microorganisms and positively control metabolism.

Delzenne and Roberfroid (1994) provided the scientific background for the characterization of such feedstuffs as pre-biotics in the sense of "*undigestible dietary ingredients which positively affect the host by beneficially and selectively stimulating the growth and/or activity of a limited number of bacteria*".

Pre-biotics act by stimulating the microorganisms which are present in the intestinal tract and not by integrating them, as is the case for probiotics. Pre- and probiotics do not exclude each other's function and can, or better, must be used simultaneously in order to obtain a powerful synergistic effect.

The concept of competitive exclusion and the intestinal microflora

From a microbiological point of view the concept of competitive exclusion implies the prevention of establishment of one microorganism into an environment because that habitat is already occupied by a competing microorganism better suited to maintain itself in that environment. The gut and the intestinal microflora represent a complex ecosystem in which several factors affect the composition of the microbial flora. Spring (1996) listed the regulatory mechanism involved in the regulation of the microbial ecology of the gut (Table 1).

A good knowledge of the composition of the GI microbial ecosystem and its control mechanism is required in order to improve the effectiveness of using commercial microbial probiotic cultures in animal production.

Chemical probiosis

Several molecules may play a pre-biotic role, promoting the development of certain microbial groups; microorganisms, in fact, do not use the energy and protein sources in the same fashion and have different needs for micronutrients and vitamins. It is therefore possible to stimulate the growth of special microbial species through the supply of certain substrates.

In ruminants, interesting results were obtained, both *in vivo* and *in vitro*, with the use of amino acids and organic acids (Masoero *et al.*, 1992, 1995) or peptides (Chen *et al.*, 1987). As the main category of probiotics, oligosaccharides have attracted a great deal of commercial attention; several types are currently produced and used as "additives" for breeding animals: fructo-oligosaccharides

(FOS), gluco-oligosaccharides (GOS), mannano-oligosaccharides (MOS), galacto-oligosaccharides (GAS), xilo-oligosaccharides (XOS). They may derive from plant origin (FOS and galacto-oligosaccharides), from enzymatic polysaccharide hydrolysis (FOS and XOS) or be re-synthesized *de novo* (FOS, GOS, GAS). Their limited inclusion in the diet (usually 0.1-0.3%) may improve weight gain, the feed conversion ratio and the health status. The size of such effects, however, is affected by numerous factors such as the type of "additive", the age of the animals, the species and the breeding conditions. In the field of monogastric nutrition, the use of fructo-oligosaccharides (FOS) gave good results in rabbits (Morisse *et al.*, 1992, 1993).

Table 1. Autogenic regulatory mechanism which affect the composition of the intestinal microbial flora (From Spring, 1996)

Regulatory mechanism	Control factors
Nutrient utilization	Competition for nutrient or growth factors Synergistic nutrient utilization
Attachment	Competition for receptor sites Stimulation of enteric cell turnover
Creation of a restrictive environment	pH Lactic acid production VFA production H ₂ S Eh Resistance to bile salts Induction of immunologic process
Productions of antimicrobial substances	NH ₃ H ₂ O ₂ Hemolysin Bacterial enzymes Bacteriophage Bacteriocins Antibiotics

The mechanism of action of such products most likely implies selective stimulation of special positive microbial clusters in the gut, such as *Bifidobacterium* (Unno *et al.*, 1993; Hirayama *et al.*, 1994; Howard *et al.*, 1995), *Bacteroides* and *Lactobacillus* (Takahashi *et al.*, 1996), *Pediococcus* spp. or *Enterococcus faecium*, but not *Salmonella typhimurium* (Oyazarbal and Corrier, 1996). Differences may, however, be found between strains of different species of the *Bifidobacterium* genus. The microorganisms of animal origin use inulin (a FOS) more efficiently than those isolated from the human gut. The same strains, however, are not capable of deriving energy from levans; it was also pointed out that the FOS which are best metabolized are those containing up to 5 fructose residues (McKellar *et al.*, 1993).

The efficacy of dietary integration with oligosaccharides is also modulated by the type of diet, being greater in hamsters fed with a diet containing a large amount of bran versus those receiving a feedstuff with greater meat proportion (Hirayama *et al.*, 1994). It is not only the type of diet which affects the response to FOS administration; 3 different strains of *Bifidobacterium* (*B. infantis* ATCC 15697, *B. adolescentis* ATCC 15703, *B. longum* ATCC 15707), showed different growth patterns in culture mediums added with 3 different types of fructane, a natural (extracted from *Helianthus tuberosus*) and two commercial ones (Yamazaki and Matsumoto, 1994). Galacto-oligosaccharides, even if to a lower extent than FOS, are another substrate which is capable of promoting the growth of *Bifidobacterium* and *Lactobacillus* (Morishita *et al.*, 1992). By administering GOS (polymerization grade from 1 to 7, with prevalence of units with 5 residues) to axenic rats, Valette *et al.* (1993) established that such molecules are resistant to intestinal digestion; if the same animals are

inoculated with human intestinal microflora, the subsequent administration of GOS does not change caecal pH, VFA production and lactic acid concentration. Changes are in fact observed in terms of the VFA profile (reduced molar percentages of butyric, isobutyric, isovalerianic acids and increased molar percentage of caproic acid) with increased production of H₂ and CH₄. Reduced production of branched chain VFA may point to a decrease in the proteolytic activity of the large intestine.

The administration of FOS increases the intestinal absorption of Ca, Mg, P (Ohta *et al.*, 1994; Ohta *et al.*, 1995a; Baba *et al.*, 1996) and Fe (Ohta *et al.*, 1995b) in rats, perhaps due to the effect of enzymatic hydrolysis performed by Lactobacilli on compounds which chelate such minerals, or by virtue of a lowering of the pH induced in the colon. The results obtained by Ohta *et al.* (1994) show that the absorption sites and mechanisms may considerably differ for these minerals.

FOS, galacto-oligosaccharides, malto-oligosaccharides (MO) and raffinose (RF) differ in their ability to stimulate mineral absorption; FOS are those that yield the best results, RF and galacto-oligosaccharides have an intermediate effect, while MO have no effect (Ohta *et al.*, 1993). Reduced lipidemia and liver lipid deposition have been observed in experimental animals following the inclusion of FOS in the diet (Otsuka and Kubo, 1995).

Positive results on reduction of pathogens intestinal colonization by feeding animals with MOS is probably due to the capacity of these sugars to bind to pathogenic organisms such as *Salmonella typhimurium* (Oyofu *et al.*, 1989) and *Escherichia coli* or stimulate the immune system. Both these mechanisms have a common feature: both the cell receptors and the antigenic determinants of several pathogenic bacteria contain mannans (Castro *et al.*, 1994; de Ruiter *et al.*, 1994; Kagaya *et al.*, 1996). Some oligomannans are deliberately included in vaccines as adjuvants which may enhance and prolong an immune response. If administered through the diet, oligomannans would compete with the corresponding intestinal receptors, which are substrates for the adhesion of the gut pathogens, thus reducing their ability to form colonies in the digestive tract epithelia. The same approach based on competitive adhesion between bacterial adhesion sites, enteric cellular receptors and molecules can be also use for lectins. A snowdrop-extracted lectin (GNA) showed, apart from high specificity for oligomannans, the ability to stop the growth of *E. coli* type I in the gut of rats (Pusztai *et al.*, 1993). This strategy may also be extended to the ability, by some sugars, to bind to bacterial toxins (Stoll *et al.*, 1980). A risk linked to the addition of this substance is represented by the possible increase in intestinal colonization by *S. typhimurium* due to the formation of bridge bindings between the bacterial cells and the enteric mucosa (Abud *et al.*, 1989; Pusztai *et al.*, 1990).

The administration of oligomannans to rats can stimulate macrophage activity (Newman, 1995); this probably accounts for the lower incidence of lung diseases found in calves receiving commercially available oligomannans (Newman *et al.*, 1993). Furthermore, their ability to stimulate the enteric receptors recognized by pathogenic bacteria allows them to bind to such bacteria, thus reducing their availability in the digestive tract (Spring *et al.*, 1996).

No information is available as to the effects that such products have on the processes which take place in the large intestine of ruminants. It is in fact most likely that such products cannot perform their action in this portion of the digestive tract, since they possibly undergo complete breakdown in the rumen. It is, among others, quite likely that a lack of fermentation energy systematically occurs in the large intestine. For this reason, the control of blood urea necessarily implies monitoring the energy availability in this part of the digestive tract.

New additives for pigs

Microbial cultures

Bacteria

An assessment of the results which can be obtained through the use of microbial cultures is made more complex by the at times contemporary use of antibiotics which adds to other variability factors such as: (i) age of the animal; (ii) diet; (iii) probiotic dose; (iv) microbial genus and species; and (v) viability and specificity of the microorganism.

An improvement in the health status is generally found together with reduced mortality, less consistent are, however, the effects on weight gain and feed conversion (Vanbelle *et al.*, 1990). The size of response is also linked to inadequate gastric production of HCl in piglets, which favours the settlement of pathogens in the gut. Similarly to acidifiers, the administration of lactic bacteria causes a drop in pH which may offset the lower acid secretion in the abomasum. It is not by chance that the best results are obtained with piglets early after weaning fed with a diet of plant origin containing no milk.

The results obtained during weaning are quite inconsistent, even if pointing to a certain efficiency for the addition of lactic bacteria (*Lactobacillus* and *Bifidobacterium*), alone or in combination with strains of *Streptococcus faecium*. Apart from the above factors which may affect efficiency, the following should also be mentioned:

(i) Use of different species from the *Lactobacillus* genus (*L. casei*, *L. bulgaricus*, *L. acidophilus*).

(ii) The fact that not all *Lactobacilli* isolated from the animal gut are capable of effectively resisting the pathogens. Hillman and Fox (1994) pointed out that only 3 out of 31 strains of *Lactobacillus*, isolated from pig stool, are in fact capable of strongly inhibiting the growth of *Escherichia coli* O149.

Apart from the ability to colonize the intestinal epithelium by adhering to it and producing lactic acid or bacteriocins, part of the probiotic effect of lactic bacteria could be due to the power of aggregation against *E. coli* shown by strains of *L. acidophilus* and *L. salivarius* (Spencer and Chesson, 1994).

The addition of *Bacillus subtilis* spores increases the number of lactobacilli while decreasing the number of coliform bacteria (Bonomi *et al.*, 1995; Kornegay and Risler, 1996; Maruta *et al.*, 1996). This effect, however, was found to be linked to the type of *B. subtilis* strain which is used (Kornegay and Risler, 1996). Maruta *et al.* (1996) also found a drop in the populations of Clostridia, Streptococci and Enterobacteriaceae. However, the changes induced in the intestinal microbial population have not been such as to significantly affect production performance (Martelli, 1992; Kornegay and Risler, 1996). Also less recent works had found less than 5% improvement in the daily weight gain (Peo, 1984; Trotters, 1984) or even negative variations (Pollman *et al.*, 1984).

By adding *Bacillus toyoi* spore to sow (300 ppm) and piglet (20 ppm) diets Gunther (1994) improved the weaned litter size (+0.55 piglet/sow), the growth (5.8-8.1%) and feed (3.72-6.20%) efficiencies.

Yeasts

Even if some studies (Bertin and Tournut, 1994; Roques *et al.*, 1994; Kornegay *et al.*, 1995) found better performance following the use of yeasts, only one case showed a positive effect on the microbial composition of the gut, with reduced number of pathogens (Roques *et al.*, 1994). In other studies (Jost *et al.*, 1993; Veum *et al.*, 1995; Bekaert *et al.*, 1996) no improvement in breeding performance was found in association with the administration of cultures of *Saccharomyces cerevisiae*. Even if showing no improvement in the digestibility of fibre fractions, Kornegay *et al.* (1995), were able to obtain an increase in the daily weight gain only in high-fiber diets containing 8% of peanut hulls. This would seem to indicate greater metabolic activity on the part of the cellulolytic microflora of the pig caecum, resulting in greater availability of volatile fatty acids to the animal.

The better growth and feed conversion obtained by Savoini *et al.* (1996) using cells of *S. cerevisiae* enriched with Cr are also due to the growth promoting action of this mineral and not only to the effect of the yeast on the intestinal microflora.

Feed induced anti secretory proteins (ASP)

Intestinal fluid secretion induced by enterotoxins may be inhibited by anti secretory proteins (ASP) (Lange and Lönnroth, 1984). These regulatory protein are synthesized in the central nervous system

(Lönnroth and Lange, 1986) and transported via the blood and bile to gut; they seem to play an important role against enteric disease (Goransson *et al.*, 1993). Sow's milk contains ASP and suckling pigs absorb it from the intestine by passive absorption (Sigfridsson *et al.*, 1995), ASP blood level decreases after weaning (Lange *et al.*, 1993) exposing piglets to enteric diseases. Supplementing creep feed with sugars, sugar alcohols and pure aminoacids can increase the production of feed induced lectines (FIL) which have the same antisecretory effects as ASP but a slightly different chemical structure (Lönnroth and Lange, 1986). Table 2 shows the results of field experiments comparing the effect of control and ASP-inducing diet on animal health and performances.

Table 2. The influence of a FIL-inducing diet in the production performance in 3 farms (modified from Goransson *et al.*, 1993)

Farm	No. of pigs	Daily weight gain (g) 0-35 d post weaning		ASP (units/ml plasma) 4 d post weaning		Post weaning diarrhoea (%)	
		C	FIL	C	FIL	C	FIL
1	40	202	247	0.42	0.87	35	10
2	54	266	325	0.79	1.05	15	4
3	325	284	380	0.76	0.94	31	2
Differences (%)			+26.6		+43.94		-80.26

The experimental diet caused an increase in ASP blood level and a reduction in clinical post weaning diarrhoea, with higher daily weight gain versus the control diet. The ASP plasma level sharply decreased in the first week after weaning, probably due to stress. On the contrary the blood FIL concentration did not fall, thus providing better protection against diarrhoea in young piglets.

Acidifiers

Supplementing the diets of weanling pigs with organic acids, such as citrate, formate, fumarate or propionate gives inconsistent responses in terms of performances. As reported by Ravindran and Kornegay (1993) citrate supplementation to the weaner piglet diets modified from -11.3 to 14.3% and from -6 to 11.1%, compared to control, the daily gain and feed conversion efficiency, respectively. The best responses were usually obtained during the first 3 weeks of age probably because of insufficient gastric production of HCl and with level of acid supplementation higher 30 g/kg of feed. Generally, improvements in feed efficiency tended to be more consistent than body weight gain improvements.

Several studies have reported consistent improvements in weight gain and feed conversion with the addition of fumaric acid to weaner diets, but other experiments have not provided any evidence of positive effects of fumarate supplementation on animal performances (Easter, 1988; Ravindran and Kornegay, 1993). Based on what these Authors have reported, also fumaric acid shows a dose-related effect similar to that of citrate, with increased efficacy above 20 g/kg supplementation. Also in this case the parameter mostly affected by treatment is feed conversion (Table 3).

Bolduan *et al.* (1988), Sweet *et al.* (1990) and Mathew *et al.* (1991) reported improved growth rates when weaner diets were supplemented with propionic acid (from 0.1 to 1%). In contrast, Giesting and Easter (1985) and Thacker *et al.* (1992) reported that the addition of 2 or 2.5% of propionate had no beneficial effect on the growth or feed conversion of weaner pigs; Giesting and Easter (1985) also found depressed feed intake possibly due to the pungent aroma of propionic acid.

Eckel *et al.* (1992) fed early-weaned pigs with 6, 12 and 18 g/kg of formic acid and obtained improved post-weaning growth (+23, 31 and 29% respectively); similar results were obtained by Eidelsburger *et al.* (1992a). A depression in feed intake was recorded with supplementation levels higher than 18 g/kg. The administration of formic acid reduces NH₃ concentration in the large intestine (Eidelsburger *et al.*, 1992a; Gedek *et al.*, 1992; Roth *et al.*, 1992b) and changes the composition of the intestinal microflora with increased availability of coliform bacteria (Gedek *et al.*, 1992a; Kirchgessner *et al.*, 1992a,b), reduction in *Bacteroides* (Gedek *et al.*, 1992a; Kirchgessner *et al.*, 1992a,b) lactic bacteria (Gedek *et al.*, 1992a) or lactic acid concentration; in contrast, increased production of acetic acid was also found (Eidelsburger *et al.*, 1992a; Roth *et al.*, 1992a).

Table 3. Summary of published data on the effects (% compared to control) of organic acids supplementation on the performance of weaner pigs (modified from Ravindran and Komegay, 1993)

Treatments		No. of trials	Daily gain	Feed intake	Gain/feed
Citric acid (g/kg)	10	5	+2.88	+4.28	+0.66
	15	5	-0.66	-2.82	+1.58
	20	4	+1.42	-3.10	+6.95
	30	4	+6.17	+7.37	+0.4
Citric acid (age of pigs, d)	7-10	4	+4.48	+9.30	-0.53
	20-21	7	+5.01	+4.54	-0.19
	25-32	9	-0.43	-4.05	+4.81
Fumaric acid (g/kg)	10	2	+2.95	-2.10	+1.15
	15	5	-0.62	+0.96	-0.16
	20	4	+5.93	-1.70	+7.50
	30	4	+4.05	-2.20	+5.88

Numerous hypothesis have been put forward concerning the mode of action of acidifiers, all linked to the effects induced by reduced gastric pH, such as increased nutrient digestibility (Eckel *et al.*, 1992; Eidelsburger *et al.*, 1992b; Kirchgessner *et al.*, 1992a,b; Thacker *et al.*, 1992) following more effective proteolytic enzyme activation. In contrast, other Authors (Bolduan *et al.*, 1988; Giesting and Easter, 1991; Mosenthin *et al.*, 1992; Gabert and Sauer, 1995) have not found any improvement in nutrient utilization.

The use of organic acid can reduce *E. coli* colonization in the gut (Bolduan *et al.*, 1988; Mathew *et al.*, 1991; Eckel *et al.*, 1992; Gedek *et al.*, 1992b; Isobe *et al.*, 1994) as well as the incidence of diarrhoea (Eidelsburger *et al.*, 1992b); other works have not pointed to any effects on enteric disorders or on a reduction in the number of intestinal coliform bacteria (Sutton *et al.*, 1991; Gedek *et al.*, 1992a; Kirchgessner *et al.*, 1992a,b; Risley *et al.*, 1992, 1993).

The intermediate organic acids of Krebs's cycle might act as energy sources inducing more efficient energy metabolism (Kirchgessner and Roth, 1982). Better efficiency of acidifying treatments could be obtained by resorting to their joint administration with probiotic bacteria (Johnson, 1992) while encapsulation with a protective matrix (Maxwell *et al.*, 1993) might turn out to be useful in adult subjects where the problem is not so much an insufficient gastric secretion of HCl but rather an adequate control of caecal fermentations.

According to Galfi and Neogràdys (1996) Na-salts of *n*-butyrate, if added in a percentage of 0.17% (on DM basis) to the diet, reduce intestinal colonization by pathogenic *E. coli* and increase the number of Lactobacilli. The proposed mode of action of *n*-butyrate is related to an ionophore-like action of this acid and not to a pH-lowering effect.

Essential oils

Some higher vegetable species (anise, origan, rosemary, pepper, celery, thyme, etc.) contain essential oils which give them aromatic properties. If added to the diet of fattening pigs, such substances considerably improve their growth pattern (Table 4).

Table 4. Performance of fattening pigs fed control or supplemented (flavours, antibiotics) diets (modified from Gunther, 1991)

Items	Negative control	Flavoured diets		Salocin control 20 mg/kg
		75 g/t	112.5 g/t	
Weight gain 70 d (kg)	69.60 ^a	73.20 ^b	75.75 ^b	75.40 ^b
Weight gain 70-120 d (kg)	35.55 ^a	37.65 ^b	39.20 ^c	37.85 ^{bc}
Daily weight gain (g)	628.3 ^a	685.4 ^b	709.6 ^c	704.6 ^{bc}
Relative weight gain (%)	100	109.1	112.9	112.1
Feed conversion ratio	3.11	2.93	2.95	2.95

a,b,c: P<0.01

Gunther suggests that such effects are due to: (i) lower amino acid oxidation; (ii) an antibiotic-like action against the intestinal microorganisms, with reduced thickness of the villi and, consequently, reduced protein metabolism of the enterocyte; (iii) increased activity of the digestive tract enzymes; and (iv) greater food intake because of improved palatability of the feedstuff.

Experimental evidence is currently available only for bactericidal action (Brud and Gora, 1990) and dry matter intake (Baidoo *et al.*, 1986; Villalba and Provenza, 1996), while the results concerning increased activity of the digestive tract enzymes are more uncertain (Ceroli and Fiorentini, pers. comm.).

New additives for poultry

The integration of FOS into diets fed to poultry inoculated with *S. typhimurium* versus controls causes a reduction in the intestinal colonization by this pathogen and also an improvement in the daily weight gain and feed conversion ratio (Choi *et al.*, 1994; Oyazarbal and Corrier, 1996). On the contrary, no effects were found on the dressing percentage and the meat fat content (Waldroup *et al.*, 1993). The contemporary use of FOS and cultures of probiotic bacteria makes it possible to establish interesting synergies in the fight against salmonellosis (Table 5). Several trials have shown a considerable reduction in the number of intestinal pathogens following the administration of mixed microbial cultures from faecal suspensions of healthy subjects both experimentally (Hinton *et al.*, 1991; Blankenship *et al.*, 1993; Kogut *et al.*, 1994; Hakkinen and Schneitz, 1996; Hume *et al.*, 1996) and in field conditions (Wierup *et al.*, 1992). Recording of such microbial populations by the health authorities implies that they have been microbiologically characterized (Stavric, 1992). This requires the identification of less complex mixtures which can compete with the pathogens. Schoeni and Wong (1994) were able to obtain markedly reduced intestinal colonization by *Campilobacter jejuni* with the use of mixtures containing *Citrobacter diversus*, *Klebsiella pneumonie* and *E. coli*.

The intestinal *S. dublin* and *E. coli* concentrations were reduced when oligomannans were added to broiler chicks diets (Spring *et al.*, 1996) (Fig. 1). The most probable mechanism involved is the saturation of bacterial intestinal-wall receptors reducing the bacterial chance of colonizing the intestine.

The contemporary administration of probiotic microorganisms and sugars (lactose, mannose, FOS) improves the antagonist action against *C. jejuni* (Table 6).

Table 5. Effect of 0.75% FOS in the feed and administration of an undefined bacterial culture (BC) on *Salmonella* colonization of 7-day-old chicks (Bailey *et al.*, 1991)

Treatment	Challenge level	% <i>Salmonella</i> positive chicks
None	10 ⁶	47.5
FOS	10 ⁶	36.4
BC	10 ⁶	43.5
FOS + BC	10 ⁶	10.5
None	10 ⁹	95.5
FOS	10 ⁹	87.0
BC	10 ⁹	60.9
FOS + BC	10 ⁹	19.0




Fig. 1. Effects of oligomannans addition in broiler chicks diets on intestinal *E. coli* and *S. dublin* concentrations.

Similarly, other Authors found an improvement in the reduction of intestinal colonization by *Salmonella* by administering lactose (5-10%) to animals who had already received CE cultures (Corrier *et al.*, 1991; Ziprin and DeLoach, 1993; Kogut *et al.*, 1994). Also some strains of *Lactobacillus* spp. were found to reduce the number of coliform bacteria in the caecum (Jin *et al.*, 1996a; Rada and Marounek, 1997) and of *Salmonella* (Jin *et al.*, 1996b); the presence of lactose in the diet enhances this protective action (Quinn *et al.*, 1995). An interesting side-effect found in conjunction with the administration of *Lactobacillus* is a reduction in the cholesterol content of the egg yolk (Haddadin *et al.*, 1996; Mohan *et al.*, 1996), due to the production of a carbohydrate which binds to cholesterol and either prevents its intestinal resorption or causes its precipitation in an insoluble form increasing its elimination through the faeces rather than its resorption by the gut. Better performances (body weight gain, feed/gain) were obtained also following the administration of *S. faecium* (Owings *et al.*, 1990) *Lactobacillus* spp. plus *B. bifidum*, plus *A. oryzae* and *Torulopsis* (Mohan *et al.*, 1996), *L. reuteri*

(England *et al.*, 1996) *Lactobacillus* spp. and *B. subtilis* (Jin *et al.*, 1996) *L. acidophilus* plus *B. subtilis* plus *S. faecium* (Chiang and Hsieh, 1995). The use of acidifiers significantly reduces intestinal colonization by *Salmonella* only if in combination with microbial cultures (Hinton *et al.*, 1991) but enhances growth in broilers (Skinner *et al.*, 1991).

Table 6. Ability of defined microbial cultures[†] (MC) and carbohydrate treatments to prevent intestinal colonization by *C. jejuni* (Modified from Schoeni and Wong, 1994)

Treatments	No. of trials	No. of chicks	% of positive chicks	Cecal pH
Control	6	37	61.6	5.7
MC	5	27	20.2*	5.8
Lactose with:				
No. MC	4	25	31.1	5.1*
MC	4	27	40.8	5.5*
Mannose with:				
No. MC	4	25	12.9*	5.7
MC	3	25	0.0*	5.8
FOS with:				
No. MC	4	25	7.7*	5.4*
MC	4	25	14.9*	5.3*

[†]*Citrobacter diversus*, *Klebsiella pneumoniae* and *E. coli*

*Significantly different (P<0.05) from the control value

Essential oils

Similarly to Gunther (1991), also Piva *et al.* (1991) found that dietary supplementation with essential oils improves the performance of broilers also in comparison with diets supplemented with virginiamycin (Table 7).

Table 7. Body weight gain of broilers fed diet with or without additives (virginiamycin or essential oils) (From Piva *et al.*, 1991 modified)

Average weight (g/head)	With virginiamycin		Without virginiamycin	
	Control	Treated [†]	Control	Treated [†]
1-29 days	964.57 ^a	968.72 ^a	965.75 ^a	991.06 ^b
29-44 days	1867.81 ^{ABb}	1981.74 ^{Bc}	1779.09 ^{Aa}	1932.39 ^{Bbc}
44-61 days	2944.09 ^{ABb}	3022.50 ^{Bbc}	2839.56 ^{Aa}	3064.35 ^{Bc}

[†]with CRINAROM 737

a,b,c: P<0.05; A,B: P<0.01

New additives for rabbits

In rabbits, the administration of FOS reduces the pH and the NH₃ concentration in the caecum, increases the production of VFA and favours the presence of non-pathogenic strains of *E. coli* (Morisse *et al.*, 1993) (Table 8).

Table 8. Effect of adding fructo-oligosaccharydes on rabbit health status, performance and caecum content (Modified from Morisse *et al.*, 1993)

Treatments	pH	<i>E. coli</i> (log ₁₀ /g)	Enteritis (%)	NH ₃ (mM/kg)	VFA (mM/kg)		
					C ₂	C ₃	C ₄
Control	6.26 ^b	2.5 ^A	46.4 ^B	17.0 ^B	45.7 ^a	1.3	9.2 ^a
Treated	6.04 ^a	4.2 ^B	14.8 ^A	11.1 ^A	57.6 ^b	1.8	19.8 ^b

a,b,c: P<0.05; A,B: P<0.01

Luicke *et al.* (1992) could not find any differences in weight gain, in the feed conversion ratio or in the composition of the faecal contents of rabbits who had previously received FOS. Dietary supplementation with microbial cultures may be considered as an alternative to the use of antibiotics.

Contemporary administration of *L. acidophilus*, *Streptococcus faecium* and *S. cerevisiae* (LACTO-SACC) increases the digestibility of the diet (Gippert *et al.*, 1992; Yamani *et al.*, 1992; Kamra *et al.*, 1996), reduces the incidence of enteric diseases (Hollister *et al.*, 1989, 1990), especially in conjunction with rations having a high starch content (Nieves-Delgado *et al.*, 1992), increases weight gain (Gippert *et al.*, 1992; Yamani *et al.*, 1992; Ayyat *et al.*, 1996) and improves feed conversion (Hollister *et al.*, 1989, 1990). Similarly to Luicke *et al.* (1992), also Kamra *et al.* (1996) found no positive change in performance in spite of increased protein digestibility.

By supplementing the diet with spores from the *Bacillus* genus, Voros and Gaal (1992) found less coliform bacteria in the gut but no effects on growth, similarly to the results obtained by Maertens *et al.* (1994) in the post-weaning phase. In the latter trial, however, a slight improvement (2.3%) was detected in the conversion ratio together with higher weight gain in the pre-weaning phase. Instead, Zoccarato *et al.* (1995) were able to improve weight gain, feed conversion ratio and nutrient digestibility by supplementing the diets with *B. subtilis* and *B. licheniformis*.

The addition of 0.15% live cells of *S. cerevisiae* to the feedstuff made it possible to reduce mortality and increase liveweight both at the time of weaning and after 70 d; at a level of 1% the results were less good (Maertens and De Groote, 1992).

The contemporary administration of 0.5 g/l of acidifiers and microorganisms (Acid Pack 4 way) increases the fermentation activity in the caecum (Kermauner and Strucklec, 1996), thus accounting for better raw fibre digestibility and improved weight gain as found by Yamani *et al.* (1992). However, a drop in performance was observed when the level of treatment was increased to 2 g/l.

New additives for ruminants

Calves

The results reported in the literature about the use of Lactobacilli are not homogeneous; several studies found no improvement in health status and performance following the administration of lactic bacteria (Jenny *et al.*, 1991; Harp *et al.*, 1996). McCormick (1984), reviewing results obtained in the USA with different strains of *Lactobacillus*, noted that only 2 of 10 experiments were positive.

Higginbotham and Bath (1993) and Cruywagen *et al.* (1996) found a non significant improvement in growth rate only in the first 2 weeks after birth in calves fed *Lactobacillus*-supplemented diets.

On the other hand some researchers obtained better performances (Abe *et al.*, 1995) and health status (Abe *et al.*, 1995; Abu-Tarbusch *et al.*, 1996) in calves following the addition of *Lactobacillus* to the diet (Table 9).

Table 9. Performance and incidence of diarrhoea in calves fed probiotics with or without antibiotics (From Abe *et al.*, 1995)

Item	Antibiotics [†] added to the starter diets			No. antibiotics added	
	<i>Bifidobacterium pseudolongum</i>	<i>Lactobacillus acidophilus</i>	Control	Probiotics	Control
No. of calves	15	15	15	10	9
Final weight (kg)	79.3	77.2	71.8	79.9	73.4
Weight gain (kg)	31.8 ^b	30.9 ^b	25.4 ^a	40.5	36.2
Feed/gain	2.1 ^a	2.07 ^a	2.37 ^b	1.59	1.64
Fecal score ^{††}	0.19	0.16	0.23	n.r.	n.r.
Diarrhoea cases	n.r.	n.r.	n.r.	1 ^a	7 ^b

[†]Colistin sulphate 20 g/t and Zn-bacitracin 4.2×10^6 units/t

^{††}Fecal score: 1 = normal; 2 = soft, 3 = scours

a,b: P<0.05

Oligosaccharides, too, have been used in calves with positive results (Webb *et al.*, 1992; Newman *et al.*, 1993; Quigley, 1996; Quigley *et al.*, 1997) (Table 10).

 Table 10. Body weight (BW) gain, incidence of scours, dry matter intake (DMI) and feed efficiency of calves fed milk replacer (MR) containing antibiotics[†] (AB) or galactosyl-lactose (GL) for 26 days (From Quigley *et al.*, 1997)

Item	Treatments			Contrast	
	MR	AB	GL	MR vs AB	MR vs GL
No. of calves	32	32	32	-	-
BW gain (g/d)	125	177	197	0.07	0.02
Fecal score ^{††}	2.27	2.07	2.06	0.07	0.08
DMI (g/d)	475	474	475	NS	NS
BW gain/DMI (g/kg)	282	399	443	0.07	0.08

[†]138 mg/kg oxytetracycline and 276 mg/kg neomycin

^{††}Fecal score: 1 normal to 4 = sever scours

On the whole, the results obtained by adding probiotics to artificial milk were indeed affected by the environmental conditions, since significant improvements could be achieved when the animals were exposed to stress (Schwab *et al.*, 1980; Seymour *et al.*, 1995), while in normal environmental conditions no improvement was found after the addition of non-conventional sugars to the diet (Jenny *et al.*, 1991; Morrill *et al.*, 1995). As previously reported, by supplementing the diets of calves with lactic acid bacteria, Abe *et al.* (1995) were able to increase performance and reduce scouring, particularly when no antibiotics were fed to the animals.

Adult ruminants

In adult ruminants, antimicrobial agents are used as growth promoters in fattening animals for the prevention of ruminal acidosis and, to a lesser extent, for the control of caecal fermentations; several active ingredients have been approved by the EU (Virginiamycin, Flavomycin, Na-Monensin). Until a few months ago, avoparcin was also allowed and its use extended to dairy cows; subsequently it was

banned because of alleged interactions with vancomycin. Such molecules mainly perform their function in the rumen by modulating fermentations and allowing the host animal to achieve greater efficiency, with increased propionate content and reduced lactic acid concentration. Lower protein degradation in the rumen and less CH₄ were also observed. These results are accompanied by other effects like reduced fibre breakdown or reduced protein synthesis in the rumen.

It is, however, possible to manipulate certain biochemical events and the microbial composition of the rumen by means of a probiotic approach. Feeding steers with rations added with cells from *S. cerevisiae* reduces the production of lactate and stabilizes the pH (Williams *et al.*, 1991); this is most likely due to a stimulating action on the bacteria which use lactic acid, e.g., *Selenomonas ruminantium* (Nisbet and Martin, 1991) or *Megasphaera elsdenii* (Rossi *et al.*, 1995); the addition of malic acid to the diet of fattening steers also stabilizes the ruminal pH and improves production performance (Streeter *et al.*, 1994), probably by stimulating *S. ruminantium* (Nisbet and Martin, 1991; Callaway and Martin, 1997) or *M. elsdenii* (Rossi *et al.*, 1995; Chaucheyras *et al.*, 1996). An increase in the use of lactate in *S. ruminantium* can be obtained by using products based on *Aspergillus oryzae* (Nisbet and Martin, 1990). Several studies have pointed to a faster breakdown of the raw fibres in rations supplemented with yeast cultures (Williams *et al.*, 1991; Carro *et al.*, 1992) or *A. oryzae* (Gómez-Alarcón *et al.*, 1990, 1991). They also found increased acetate concentrations in the rumen (Piva *et al.*, 1993), probably due to increased growth rate among the cellulolytic microflora (Dawson *et al.*, 1990; Callaway and Martin, 1997).

According to Yoon and Stern (1995) (Table 11), the administration of *S. cerevisiae* or *A. oryzae* causes an appreciable improvement in the growth of calves and steers.

Table 11. Effects of *Saccharomyces cerevisiae* or *Aspergillus oryzae* on performances of growing cattles (variation compared to control) (Modified from Yoon and Stern, 1995)

Animal	Dosage (g/d)	Positive trials on total studies	Dry matter intake (%)	Weight gain (g/d)	Feed/gain (%)
<i>S. cerevisiae</i>					
Calves	28.6 ± 51.3	7/8	+6.2	+93.7 (12.1%)	-6.2
Bulls	9.2 ± 1.2	2/2	+4.3	+30.0 (1.9%)	-2.4
Steers	109.5 ± 140.7	2/2	+6.4	+105.0 (8.04%)	-6.0
<i>A. oryzae</i>					
Steers	90 - 113	2/2	n.a.	+45 (4.8%)	n.a.
Calves	0.5 - 3	2/2	+5	+55 (10.5%)	-4.61

n.a.: data not available

The ruminal bacteria have different nutritional requirements in terms of energy and nitrogen (Cotta, 1993; Stewart and Bryant, 1988), peptides (Chen *et al.*, 1987), amino acids and organic acids (Russell and Sniffen, 1984; Masoero *et al.*, 1992). The use of amino acids and organic acids changes the ruminal fermentation profile and microbial growth both *in vitro* (Stack and Cotta, 1986; Masoero *et al.*, 1992; Crepaldi *et al.*, 1994; Barbour *et al.*, 1996) and *in vivo*, with positive effects on the use of feedstuffs and animal weight gain (Masoero *et al.*, 1995; Streeter *et al.*, 1994). In sheep, lactose supplementation in a ration containing corn silage caused an increase in the duodenal flow of microbial proteins (Chamberlain *et al.*, 1993). It is therefore likely that the growth of some microbial groups can be increased to the detriment of others by adequately administering sugars, amino acids or organic acids. Morvan and Fonty (1995) found that the addition of xylose promotes the growth of an acetogenic bacterium which is capable of producing acetate starting from H₂ and CO₂, which would reduce CH₄ production without resorting to antibiotics; similar results were obtained by Chaucheyras *et al.* (1995) using yeast cultures.

Conclusions

The concerns about the possible effects of antimicrobial agents used as growth promoters in animal breeding in terms of increasing antibiotic resistance towards both human and animal pathogens, requires the adoption of an entirely different approach to growth promoters.

Several practical possibilities have been advanced: from rationalizing the use of pro- and prebiotics to adopting nutritional strategies which adequately enhance the immunity and control features of the secretory factors normally found in feedstuffs. The rations shall have to be designed so as to feature precise pharmacological characteristics in order to help minimize the use of conventional drugs in the prevention and treatment of diseases.

Futures developments

Ruminants

Use of rumen-protected oligosaccharides (fructans and mannans) to modulate the microbial activity of the gut. In the rumen, the administration of non-conventional sugars or oligosaccharides could play a certain role on two occasions:

- (i) Increased competitiveness between genetically modified ruminal microorganisms.
- (ii) Increased competitiveness between important ruminal microorganisms which are usually present in small numbers and administered orally.
- (iii) Stimulation of the growth of microorganisms which occupy special ecological niches such as the acetogenic bacteria which use H_2 and reduce CH_4 production. Such bacteria may use xylose, a sugar which is used relatively little by the dominant microbial species. The reduction in CH_4 emissions is an important objective in controlling the environmental impact of animal breeding farms.

Also the administration of trace elements like Cu and Zn, in highly available forms (chelates), adequately protected against ruminal fermentation and performing an antibacterial (Cu), immunostimulating (Zn) or antioxidating (Se) action, might significantly contribute to the modulation of intestinal fermentations.

Monogastrics

Special carbohydrates should be used which selectively stimulate the lactic microflora available in the gut in order to limit both putrefaction and the release of toxic agents.

- (i) New intestinal microorganisms should be isolated which can effectively compete with pathogens.
- (ii) The polymerization degree of different oligosaccharides should be studied in order to assess the most effective one in the stimulation of the autochthonous intestinal bacteria.
- (iii) The probiotic effect of lectins derived from garlic, banana and shallot should be further investigated although interesting results have already been obtained (Koshte *et al.*, 1990; Kaku *et al.*, 1992; Mo *et al.*, 1993).
- (iv) Study of other molecules like L-fucose, N-acetyl-D-glucosamine and N-acetyl-D-neuraminic acid able of preventing intestinal walls pathogens adhesion.

References

Abe, F., Ishibashi, N. and Shimamura, S. (1995). Effect of administration of bifidobacteria and lactic acid bacteria to newborn calves and piglets. *J. Dairy Sci.*, 78: 2838-2846.

- Abud, R.L., Lindquist, B.L., Ernst, R.K., Merick, J.M., Lebenthal, E. and Lee, P.C. (1989). Concanavalin A promotes adherence of *Salmonella typhimurium* to small intestinal mucosa of rats. *Proc. Soc. Exp. Biol. Med.*, 192: 81-86.
- Abu-Tarboush, H.M., Al-Saiady, M.Y. and Keir El Din, A.H. (1996). Evaluation of diet containing lactobacilli on performance, fecal coliform and lactobacilli of young dairy calves. *Anim. Feed Sci. Technol.*, 57: 39-49.
- Ayyat, M.S., Marai, I.F.M. and El-Aasar, T.A. (1996). New Zealand white rabbit does and their growing offsprings as affected by diets containing different protein level with or without Lacto-Sacc supplementation. *World Rabbit Sci.*, 4: 225-230.
- Baba, S., Ohta, A., Ohtsuki, M., Takizawa, T., Adachi, T. and Hara, H. (1996). Fructooligosaccharides stimulate the absorption of magnesium from the hindgut in rats. *Nutr. Res.*, 16: 657-666.
- Baidoo, S.K., McIntosh, M.K. and Aherne, F.X. (1986). Selection preference of starter pigs fed canola meal supplemented diets. *Can. J. Anim. Sci.*, 66: 1039-1049.
- Bailey, J.S., Blankenship, L.C. and Cox, N.A. (1991). Effect of fructooligosaccharide on *Salmonella* colonization of the chicken intestine. *Poultry Sci.*, 70: 2433-2438.
- Barbour, A.F., Solaiman, S.G., Mackie, R.I., Maloney, M.A. and Murphy, M.R. (1996). The effect of stereoisomer of aspartic acid and lysine on rumen fermentation of alfalfa and bermudagrass hay *in vitro*. *J. Anim. Sci.*, 74(Suppl. 1): 285 (Abstr. 713).
- Blankenship, L.C., Bailey, J.S., Cox, N.A., Stern, N.J., Brewer, R. and Williams, O. (1993). Two-step mucosal competitive exclusion flora treatment to diminish *Salmonellae* in commercial broiler chickens. *Poultry Sci.*, 72: 1667-1672.
- Bolduan, G., Jung, H., Schneider, R., Block, J. and Klenke, B. (1988). Effect of propionic and formic acid in piglets. *J. Anim. Physiol. Anim. Nutr.*, 59: 72-78.
- Brud, W.S. and Gora, J. (1990). Biological activity of essential oils and its possible applications. In: *Proc. XI Int. Congress of Essential oils, Fragrance and Flavours*, New Delhi, 1989, Bhattacharyya, S.C., Sen, N. and Sethi, K.L. (eds). Aspect Publishing, London, pp. 13-23.
- Callaway, E.S. and Martin, S.A. (1997). Effects of a *Saccharomyces cerevisiae* culture on ruminal bacteria that utilize lactate and digest cellulose. *J. Dairy Sci.*, 80: 2035-2044.
- Carro, M.D., Lebzien, P. and Rohr, K. (1992). Effects of yeast culture on rumen fermentation, digestibility and duodenal flow in dairy cows fed a silage based diet. *Livest. Prod. Sci.*, 32: 219-229.
- Castro, M., Ralston, N.V.C., Morgenthaler, T.I., Rohrbach, M.S. and Limper, A.H. (1994). *Candida albicans* stimulates arachidonic acid liberation from alveolar macrophages through α -mannan and β -glucan cell wall components. *Infect. Immun.*, 62: 3138-3145.
- Chamberlain, D.G., Robertson, S. and Choung, J.J. (1993). Sugars versus starch as supplements to grass silage: effects on ruminal fermentation and the supply of microbial protein to the small intestine, estimated from the urinary excretion of purine derivatives, in sheep. *J. Sci. Food Agr.*, 63: 189-194.
- Chaucheyras, F., Fonty, G., Bertin, G., Salmon, J. and Gouet, P. (1996). Effects of a strain of *Saccharomyces cerevisiae* (Levucell SC), a microbial additive for ruminants on lactate metabolism *in vitro*. *Can. J. Microbiol.*, 42: 927-933.
- Chaucheyras, F., Fonty, G., Bertin, G. and Gouet, P. (1995). *In vitro* H₂ utilization by a ruminal acetogenic bacterium cultivated alone or in association with an archaea methanogen is stimulated by a probiotic strain of *Saccharomyces cerevisiae*. *Appl. Environ. Microbiol.*, 61: 3466-3467.

- Chen, G., Strobel, H.J., Russell, J.B. and Sniffen, C.J. (1987). Effect of hydrophobicity on utilization of peptides by ruminal bacteria *in vitro*. *Appl. Environ. Microbiol.*, 53: 2021-2025.
- Chiang, S.H. and Hsieh, W.M. (1995). Effect of direct-fed microorganisms on broiler growth performance and litter ammonia level. *Asian Austr. J. Anim. Sci.*, 8: 159-162.
- Choi, K.H., Namkung, H. and Paik, I.K. (1994). Effects of dietary fructooligosaccharides on the suppression of intestinal colonization of *Salmonella typhimurium* in broilers chickens. *Korean J. Anim. Sci.*, 36: 271-284.
- Corrier, D.E. Jr., Hinton, A., Kubena, L.F., Ziprin, R.L. and DeLoach, J.R. (1991). Decreased *Salmonella* colonization in turkey poults inoculated with anaerobic cecal microflora and provided dietary lactose. *Poultry Sci.*, 70: 1345-1350.
- Cotta, M.A. (1993). Utilization of xylooligosaccharides by selected ruminal bacteria. *Appl. Environ. Microbiol.*, 59: 3557-3563.
- Coussement, P. (1995). Pre- and synbiotics with inulin and oligofructose. Promising developments in functional foods. *Food Technol. Europe*, 2: 102-104.
- Crepaldi, P., Zanchi, R., Pacini, N., Moller, F. and Cicogna, M. (1994). *In vitro* effects of orotic acid on rumen flora. *Ann. Microbiol. Enzimol.*, 44: 317-326.
- Cruywagen, C.W., Jordaan, J. and Venter, L. (1996). Effects of *Lactobacillus acidophilus* supplementation of milk replacer on preweaning performance of calves. *J. Dairy Sci.*, 79: 483-486.
- Dawson, K.A., Newman, K.E. and Boling, J.A. (1990). Effects of microbial supplements containing yeast and lactobacilli on roughage-fed ruminal microbial activities. *J. Anim. Sci.*, 68: 3392-3398.
- de Ruiter, G.A., van der Bruggen, A.W., Mischnick, P., Smid, P., van Boom, J.H., Notermans, S.H.W. and Rombouts, F.M. (1994). 2-O-methyl-D-mannose residues are immunodominant in extracellular polysaccharides of *Mucor racemosus* and related molds. *J. Biol. Chem.*, 269: 4299-4306.
- Delzenne, N.M. and Roberfroid, M.R. (1994). Physiological effects of non-digestible oligosaccharides. *Lebensmittel Wissenschaft Technol.*, 27: 1-6.
- Easter, R.A. (1988). Acidification of diets for pigs. In: *Recent advances in animal nutrition*. Haresign, W. and Cole, D.J.A. (eds). Butterworth, London, UK, pp. 61-72.
- Eckel, B., Kirchgessner, M. and Roth, F.X. (1992a). Influence of formic acid on daily weight gain, feed intake, feed conversion rate and digestibility. 1. Communication: Investigations about the nutritive efficacy of organic acids in the rearing of piglets. *J. Anim. Physiol. Anim. Nutr.*, 67: 93-100.
- Eckel, B., Roth, F.X., Kirchgessner, M. and Eidelsburger, U. (1992b). Influence of formic acid on concentrations of ammonia and biogenic amines in the gastrointestinal tract. 4. Investigations about the nutritive efficacy of organic acids in the rearing of piglets. *J. Anim. Physiol. Anim. Nutr.*, 67: 198-205.
- Eidelsburger, U., Kirchgessner, M. and Roth, F.X. (1992a). Influence of formic acid, calcium formate and sodium hydrogen carbonate on dry matter content, pH value, concentration of carboxylic acids and ammonia in different segments of the gastrointestinal tract. 8. Investigations about the nutritive efficacy of organic acids in the rearing of piglets. *J. Anim. Physiol. Anim. Nutr.*, 68: 20-32.
- Eidelsburger, U., Kirchgessner, M. and Roth, F.X. (1992b). Influence of fumaric acid, hydrochloric acid, sodium formate, tylosin and toyocerin on daily weight gain, feed intake, feed conversion rate and digestibility. 11. Investigations about the nutritive efficacy of organic acids in the rearing of piglets. *J. Anim. Physiol. Anim. Nutr.*, 68: 82-92.

- England, J.A., Watkins, S.E., Saleh, E., Waldroup, P.W., Casas, I. and Burnham, D. (1996). Effects of *Lactobacillus reuteri* on live performance and intestinal development of male turkeys. *J. Appl. Poultry Res.*, 5: 311-324.
- Gabert, V.M. and Sauer, W.C. (1995). The effect of fumaric acid and sodium fumarate supplementation to diets for weanlings pigs on amino acid digestibility and volatile fatty acid concentrations in ileal digesta. *Anim. Feed Sci. Technol.*, 53: 243-254.
- Galfi, P. and Neogràdy, S. (1996). Short chain fatty acids (acidifiers) as probiotics in diets for piglets. In: *Proc. IV Int. Feed Production Conference*, Piacenza, Piva, G. (ed.). pp. 185-204.
- Gedek, B., Kirchgessner, M., Eidelsburger, U., Wiehler, S., Bott, A. and Roth, F.X. (1992a). Influence of formic acid on the microflora in different segments of the gastrointestinal tract. 5. Investigations about the nutritive efficacy of organic acids in the rearing of piglets. *J. Anim. Physiol. Anim. Nutr.*, 67: 206-214.
- Gedek, B., Roth, F.X., Kirchgessner, M., Wiehler, S., Bott, A. and Eidelsburger, U. (1992b). Influence of formic acid hydrochloric acid, sodium formate, tylosin and toyocerin on the microflora in different segments of the gastrointestinal tract. 14. Investigations about the nutritive efficacy of organic acids in the rearing of piglets. *J. Anim. Physiol. Anim. Nutr.*, 68: 209-217.
- Giesting, D.W. and Easter, R.A. (1985). Response of starter pigs to supplementation of corn soybean meal diets with organic acids. *J. Anim. Sci.*, 60: 1288-1294.
- Giesting, D.W. and Easter, R.A. (1991). Effect of protein source and fumaric acid supplementation on apparent ileal digestibility of nutrient by young pigs. *J. Anim. Sci.*, 69: 2497-2503.
- Gippert, T., Virag, G. and Nagy, I. (1992). Lacto-Sacc in rabbit nutrition. *J. Appl. Rabbit Res.*, 15:B 1101-1104.
- Gómez-Alarcón, R.A., Dudas, C. and Huber, J.T. (1990). Influence of *Aspergillus oryzae* on rumen and total tract digestibility of dietary components. *J. Dairy Sci.*, 73: 703-710.
- Gómez-Alarcón, R.A., Huber, J.T., Higginbotham, G.E., Wiersma, F., Ammon, D. and Taylor, B. (1991). Influence of feeding *Aspergillus oryzae* fermentation extract on the milk yields, eating patterns and body temperature of lactating cows. *J. Dairy Sci.*, 69: 1733-1740.
- Goransson, L., Martinsson, K., Lange, S. and Lönnroth, I. (1993). Feed-induced lectines in piglets. *J. Vet. Med. Series B*, 40: 478-484.
- Gunther, K.D. (1991). Spices in animal feed. *The home mixer*, (2).
- Gunther, K.D. (1994). The role of probiotics as feed additives in animal nutrition. In: *Proc. IV Int. Feed Production Conference*, Piacenza, Piva G. (ed.). pp. 97-114.
- Haddadin, M.S.Y., Abdulrahim, S.M., Hashlamoun, E.A.R. and Robinson, R.K. (1996). The effect of *Lactobacillus acidophilus* on the production and chemical composition of hen's eggs. *Poultry Sci.*, 75: 491-494.
- Hakkinen, M. and Schneitz, C. (1996). Efficacy of a commercial competitive exclusion product against a chicken pathogenic *Escherichia coli* and *E. coli* O157:H7. *Vet. Rec.*, 139: 139-141.
- Harp, J.A., Jardon, P., Atwill, E.R., Zylstra, M., Checél, S., Goff, J.P. and De Simone, C. (1996). Field testing of prophylactic measures against *Cryptosporidium parvum* infection in calves in a California dairy herd. *Am. J. Vet. Res.*, 57: 1586-1588.
- Higginbotham, G.E. and Bath, D.L. (1993). Evaluation of *Lactobacillus* fermentation cultures in calf feeding system. *J. Dairy Sci.*, 76: 615-620.

- Hillman, K. and Fox, A. (1994). Effect of porcine faecal lactobacilli on the rate of growth of enterotoxigenic *Escherichia coli* O149:K88:K 91. *Lett. Appl. Microbiol.*, 19: 497-500.
- Hinton, M., Mead, G.C. and Impey, C.S. (1991). Protection of chicks against environmental challenge with *Salmonella enteritidis* by competitive exclusion. *Lett. Appl. Microbiol.*, 12: 69-71.
- Hollister, A.G., Cheeke, P.R., Robinson, K.L. and Patton, N.M. (1989). Effect of water-administered probiotics and acidifiers on growth, feed conversion and enteritis mortality of weanling rabbits. *J. Appl. Rabbit Res.*, 12: 143-147.
- Hollister, A.G., Cheeke, P.R., Robinson, K.L. and Patton, N.M. (1990). Effect of dietary probiotics and acidifiers on performance of weanling rabbits. *J. Appl. Rabbit Res.*, 13: 6-9.
- Howard, M.D., Gordon, D.T., Pace, L.W., Garleb, K.A. and Kerley, M.S. (1995). Effects of dietary supplementation with fructooligosaccharides on colonic microbiota populations and epithelial cell proliferation in neonatal pigs. *J. Pediat. Gastroenterol. Nutr.*, 21: 297-303.
- Hume, M.E., Corrier, D.E., Nisbet, D.J. and DeLoach, J.R. (1996). Reduction of *Salmonella* crop and cecal colonization by a characterized competitive exclusion culture in broilers during grow-out. *J. Food Protect.*, 59: 688-693.
- Isobe, Y., Shibata, F., Komaki, H. and Kamada, A. (1994). Influence of fumaric acid administration on intestinal microorganism in weanling pigs. *Anim. Sci. Technol.*, 65: 59-66.
- Jenny, B.F., Vandijk, H.J. and Collins, J.A. (1991). Performance and fecal flora of calves fed a *Bacillus subtilis* concentrate. *J. Dairy Sci.*, 74: 1968-1973.
- Jin, L.Z., Ho, Y.W., Abdullah, N. and Jalaludin, S. (1996a). Influence of dried *Bacillus subtilis* and lactobacilli cultures on intestinal microflora and performance in broilers. *Asian Austr. J. Anim. Sci.*, 9: 397-403.
- Jin, L.Z., Ho, Y.W., Abdullah, N. and Jalaludin, S. (1996b). Antagonistic effects of intestinal *Lactobacillus* isolates on pathogens of chicken. *Lett. Appl. Microbiol.*, 23: 67-71.
- Johnson, R. (1992). Acidification of pig diets to improve health and growth of weaners. *Feed Compounder*, 12: 44-48.
- Kagaya, K., Miyakawa, Y., Suzuki, M., Nakase, T. and Fukazawa, Y. (1996). Role of mannan on adhesion of *Candida albicans* to epithelial cells. *Jpn. J. Med. Mycol.*, 37: 71-76.
- Kaku, H., Goldstein, I.J., van Damme, E.J.M. and Peumans, W.J. (1992). New mannose-specific lectin from garlic (*Alium sativum*) and ramsons (*Alium ursinum*) bulbs. *Carbohydr. Res.*, 229: 347-353.
- Kamra, D.N., Chaudhary, L.C., Singh, R. and Pathak, N.N. (1996). Influence of feeding probiotics on growth performance and nutrient digestibility in rabbits. *World Rabbit Sci.*, 4: 85-88.
- Kermauner, A. and Struklec, M. (1996). Addition of probiotics to feeds with different energy and ADF content in rabbits. 1. Effects on the digestive organs. *World Rabbit Sci.*, 4: 187-193.
- Kirchgessner, M., Eckel, B., Roth, F.X. and Eidelsburger, U. (1992a). Influence of formic acid on carcass composition and retention of nutrients. 2. Investigations about the nutritive efficacy of organic acids in the rearing of piglets. *J. Anim. Physiol. Anim. Nutr.*, 68: 73-81.
- Kirchgessner, M., Gedek, B., Wiehler, S., Bott, A., Eidelsburger, U. and Roth, F.X. (1992b). Influence of formic acid, calcium formate and sodium hydrogen carbonate on the microflora in different segments of the gastrointestinal tract. 10. Investigations about the nutritive efficacy of organic acids in the rearing of piglets. *J. Anim. Physiol. Anim. Nutr.*, 68: 73-81.

- Kirchgessner, M. and Roth, F.X. (1982). Fumaric acid as a feed additive in pig nutrition. *Pigs News Inf.*, 3: 259-264.
- Kogut, M.H., Fukata, T., Tellez, G., Hargis, B.M., Corrier, D.E. and De Loach, J.R. (1994). Effect of *Eimeria tenella* infection on resistance to *Salmonella typhimurium* colonization in broiler chicks inoculated with anaerobic cecal flora and fed dietary lactose. *Avian Dis.*, 38: 59-64.
- Koshte, V.L., van Dijk, W., van der Stelt, M.E. and Aalberse, R.C. (1990). Isolation and characterization of BanLec-1 a mannoside-binding lectin from *Musa paradisiac* (banana). *Biochem. J.*, 272: 721-726.
- Lange, S. and Lönnroth, I. (1984). Passive transfer of protection against cholera toxin in rat intestine. *Microbiol. Lett.*, 24: 165-168.
- Lönnroth, I. and Lange, S. (1986). Purification and characterization of anti secretory factor a protein in the control nervous system and in the gut which inhibits intestinal hyper secretion induced by cholera toxin. *Biochem. Biophysic. Acta*, 883: 138-144.
- Luicke, B.R., El-Sayaad, G.A.E. and Cheeke, P.R. (1992). Effects of fructooligosaccharides and yeast culture on growth performance of rabbits. *J. Appl. Rabbit Res.*, 15: 1121-1128.
- Maertens, L. and De Groote, G. (1992). Effect of dietary supplementation of live yeast on the zootechnical performances of does and weanling rabbits. *J. Appl. Rabbit Res.*, 15:B, 1079-1086.
- Maertens, L., van Renterghem, R. and De Groote, G. (1994). Effect of dietary inclusion of Paciflor R (*Bacillus* CIP 5832) on the milk composition and performances of does and on caecal and growth parameters of their weanlings. *World. Rabbit Sci.*, 2: 67-73.
- Masoero, F., Rossi, F. and Mancini, V. (1995). Effetto dell'acido aspartico sulla degradazione ruminale di fieno di prato stabile e farina di mais. In: *Atti XI Congr. Naz. ASPA*, 19-22 June, pp. 225-226.
- Masoero, F., Rossi, F. and Piva, G. (1992). Effects of organic acids, amino acids and isoacids on volatile fatty acids, bacterial dry matter production and degradation of cellulose by rumen bacteria. *In vitro* trials. *Microbiol. Alim. Nutr.*, 10: 141-146.
- Mathew, A.G., Sutton, A.L., Scheidt, A.B., Forsyth, D.M., Patterson, J.A. and Kelly, D.T. (1991). Effects of a propionic acid containing additive on performance and intestinal microbial fermentation of the weanling pigs. In: *Proc. 6th Int. Symp. on the Digestive Physiology in pigs*, PUDOC, Wageningen, The Netherlands, pp. 464-469.
- Maxwell, C.V., Sohn, K.S. and Brock, K.S. (1993). Effect of acidification on starter pig performance. *Anim. Sci. Res. Rep.*, Oklahoma State Univ., USA, pp. 333-339.
- McCormick, M.E. (1984). Probiotics in ruminant nutrition and health. In: *Proc. Georgia Nutr. Conference for the feed industry*, pp. 62-69.
- McKellar, R.C., Modler, H.W. and Mullin, J. (1993). Characterization of growth and inulase production by *Bifidobacterium* spp. on fructooligosaccharides. *Bifidobacteria and microflora*, 12: 75-86.
- Mo, H.Q., van Damme, E.J.M., Peumans, W.J. and Goldstein, I.J. (1993). Purification and characterization of a mannose-specific lectin from shallot (*Allium ascalonicum*) bulbs. *Arch. Biochem.*, 306: 431-438.
- Mohan, B., Kadirvel, R., Natarajan, A. and Bhaskaran, M. (1996). Effect of probiotic supplementation on growth, nitrogen utilization and serum cholesterol in broilers. *Brit. Poultry Sci.*, 37: 395-401.
- Morishita, Y., Konishi, Y., Tanaka, R. and Mutai, Y. (1992). The effect of transgalactosylated oligosaccharide (TOS) on the intestinal microflora and cecal short-chain fatty acids in mice and rats. *Bifidus*, 6: 11-17.

- Morisse, J.P., Maurice, R., Boilletot, E. and Cotte, J.P. (1992). Effect of a fructooligosaccharides compounds in rabbit experimentally infected with *E. coli* O.103. *J. Appl. Rabbit Res.*, 15: 1137-1143.
- Morisse, J.P., Maurice, R., Boilletot, E. and Cotte, J.P. (1993). Assessment of the activity of a fructo-oligosaccharides on different caecal parameters in rabbit experimentally infected with *E. coli* O.103. *Ann. Zootech.*, 42: 81-87.
- Morvan, B. and Fonty, G. (1995). Mixotrophy by rumen acetogenic bacteria in the utilization of hydrogen and sugars. In: *Satellite Symp. on Rumen Microbiology*, Clermont Ferrand, 16-17 September 1995.
- Mosenthin, R., Sauer, W.C., Ahrens, F., De Lange, C.F.M. and Bornholdt, U. (1992). Effect of dietary supplements of propionic acid, siliceous earth or a combination of these on the energy, protein and amino acid digestibilities and concentration of microbial metabolites in the digestive tract of growing pigs. *Anim. Feed Sci. Technol.*, 37: 245-255.
- Newman, K.E., Jacques, K. and Buede, R.P. (1993). Effect of mannanoligosaccharide on performance of calves fed acidified and nonacidified milk replacers. *J. Anim. Sci.*, 71(Suppl. 1): 271 (Abstr.).
- Newman, K.E. (1995). Oligosaccaridi mannani e nutrizione animale. In: *9th Alltech European Lectures Tour*, pp. 51-56.
- Nieves-Delgado, D., Pro-Martínez, A., Herrera-Haro, J. and Velázquez, J. (1992). Effect of probiotics on the use of mash diets with high and low alfalfa meal content in rabbits. *J. Appl. Rabbit Res.*, 15:B, 1160-1162.
- Nisbet, D.J. and Martin, S.A. (1990). Effect of dicarboxylic acids and *Aspergillus oryzae* fermentation extract on lactate uptake by the ruminal bacterium *Selenomonas ruminantium*. *Appl. Environ. Microbiol.*, 56: 3515-3518.
- Nisbet, D.J. and Martin, S.A. (1991). Effect of a *Saccharomyces cerevisiae* culture on lactate utilization by the ruminal bacterium *Selenomonas ruminantium*. *J. Anim. Sci.*, 69: 4628-4633.
- Ohta, A., Ohtsuki, M., Baba, S., Adachi, T., Sakata, T. and Sakaguchi, E. (1995a). Calcium and magnesium absorption from the colon and rectum are increased in rats fed fructooligosaccharides. *J. Nutr.*, 125: 2417-2424.
- Ohta, A., Ohtsuki, M., Baba, S., Takizawa, T., Adachi, T. and Kimura, S. (1995b). Effects of fructooligosaccharides on the absorption of iron, calcium and magnesium in iron-deficient anemic rats. *J. Nutr. Sci. Vitaminol.*, 41: 281-291.
- Ohta, A., Ohtsuki, M., Takizawa, T., Inaba, H., Adachi, T. and Kimura, S. (1994). Effects of fructooligosaccharides on the absorption of magnesium and calcium by cecectomized rats. *Int. J. Vitamin Nutr. Res.*, 64: 316-323.
- Ohta, A., Osakabe, N., Yamada, K., Saito, Y. and Hidaka, H. (1993). Effects of fructooligosaccharides and other saccharides on Ca, Mg, and P absorption in rats. *J. Jap. Soc. Nutr. Food Sci.*, 46: 123-129.
- Owings, W.J., Reynolds, D.L., Hasiak, R.J. and Ferket, P.R. (1990). Influence of dietary supplementation with *Streptococcus faecium* M-74 on broiler body weight, feed conversion, carcass characteristics and intestinal microbial colonization. *Poultry Sci.*, 69: 1257-1264.
- Oyazarbal, O.A. and Conner, D.E. (1996). Application of direct-fed microbial bacteria and fructooligosaccharodes for *Salmonella* control in broilers during feed withdrawal. *Poultry Sci.*, 75: 186-190.

- Oyofa, B.A., DeLoach, J.R., Corrier, D.E., Norman, J.O., Ziprin, R.L. and Mollenhauer, H.H. (1989). Prevention of *Salmonella typhimurium* colonization of broilers with D-mannose. *Poultry Sci.*, 68: 1357-1360.
- Piva, G., Belladonna, S., Fusconi, G. and Sicbaldi, F. (1993). Effects of yeasts on dairy cow performance, ruminal fermentation, blood components and milk manufacturing properties. *J. Dairy Sci.*, 76: 2717-2722.
- Piva, G., Morlacchini, M., Riccardi, R., Masoero, F., Prandini, A. and Mandolini, F. (1991). Probiotic effect of essential oils in animal feeding. *Microbiol. Alim. Nutr.*, 9: 161-169.
- Pusztai, A., Grant, G., King, T.P. and Clarke, M.W. (1990). Chemical probiosis. In: *Recent advances in animal nutrition*, Nottingham, 1989, Haresign, W. and Cole, D.J.A. (eds). Butterworths, pp. 47-60.
- Pusztai, A., Grant, G., Spencer, R.J., Duguid, T.J., Brown, D.S., Ewen, S.W.B., Peumans, W.J., van Damme, E.J.M. and Bardocz, S. (1993). Kidney bean lectin-induced *Escherichia coli* overgrowth in the small intestine is blocked by GNA a mannose-specific lectin. *J. Appl. Bacteriol.*, 75: 360-368.
- Quigley, J.D. III (1996). Intake, growth and health of dairy calves in response to mannanoligosaccharide and oral challenge with *Escherichia coli*. *J. Dairy Sci.*, 79(Suppl. 1): 230 (Abstr.).
- Quigley, J.D. III, Drewry, J.J., Murray, L.M. and Ivey, S.J. (1997). Body weight gain, feed efficiency and fecal scores of dairy calves in response to galactosyl-lactose or antibiotics in milk replacers. *J. Dairy Sci.*, 80: 1751-1754.
- Quin, Z.R., Fukata, T., Baba, E. and Arakawa, A. (1995). Effect of lactose and *Lactobacillus acidophilus* on the colonization of *Salmonella enteritidis* in chicks concurrently infected with *Eimeria tenella*. *Avian Dis.*, 39: 548-553.
- Rada, V. and Marounek, M. (1997). Effect of maduramicin and monensin on survival of *Lactobacillus salivarius* 51R administered in the crop and caeca of young chickens. *Arch. Anim. Nutr.*, 50: 25-29.
- Ravindran, V. and Kornegay, E.T. (1993). Acidification of weaner pig diets: a review. *J. Sci. Food Agr.*, 62: 313-322.
- Risley, C.R., Kornegay, E.T., Lindemann, M.D., Wood, C.M. and Eigel, W.N. (1992). Effect of feeding organic acids on selected intestinal content measurements at varying times postweaning in pigs. *J. Anim. Sci.*, 70: 196-206.
- Risley, C.R., Kornegay, E.T., Lindemann, M.D., Wood, C.M. and Eigel, W.N. (1993). Effect of feeding organic acids on gastrointestinal digesta measurements at various times postweaning in pigs challenged with enterotoxigenic *Escherichia coli*. *Can. J. Anim. Sci.*, 73: 931-940.
- Roth, F.X., Eckel, B., Kirchgessner, M. and Eidelsburger, U. (1992a). Influence of formic acid on pH value, dry matter content, concentration of volatile fatty acids and lactic acid in gastrointestinal tract. 3. Investigations about the nutritive efficacy of organic acids in the rearing of piglets. *J. Anim. Physiol. Anim. Nutr.*, 67: 148-156.
- Roth, F.X., Eidelsburger, U. and Kirchgessner, M. (1992b). Influence of fumaric acid, hydrochloric acid, sodium formate, tylosin and toyocerin on dry matter content, pH value, concentration of carbonic acids and ammonia in different segments of the gastrointestinal tract. 12. Investigations about the nutritive efficacy of organic acids in the rearing of piglets. *J. Anim. Physiol. Anim. Nutr.*, 68: 93-103.

- Russell, J.B. and Sniffen, C.J. (1984). Effect of C₄ and C₅ volatile fatty acids on the growth of mixed rumen bacteria *in vitro*. *J. Dairy Sci.*, 67: 987-994.
- Schoeni, J.L. and Wong, A.C.L. (1994). Inhibition of *Campylobacter jejuni* colonization in chicks by defined competitive exclusion bacteria. *Appl. Environ. Microbiol.*, 60: 1191-1197.
- Sigfridson, K., Lange, S. and Lönnroth, I. (1995). Antisecretory protein and feed induced lectines in sow and suckling piglets. In: 46th Annual Meeting of EAAP, Prague, Czech Republic (Abstr.).
- Skinner, J.T., Izat, A.L. and Waldroup, P.W. (1991). Research note: fumaric acid enhances performance of broiler chickens. *Poultry Sci.*, 70: 1444-1447.
- Spencer, R.J. and Chesson, A. (1994). The effect of *Lactobacillus* spp. on the attachment of enterotoxigenic *Escherichia coli* to isolated porcine enterocytes. *J. Appl. Bacteriol.*, 77: 215-220.
- Spring, P. (1996). *Effects of mannanoligosaccharide on different cecal parameters and on cecal concentrations of enteric pathogens in poultry*. PhD Thesis, Swiss Fed. Inst. Technology, Zurich.
- Spring, P., Dawson, K.A., Newman, K.E. and Wenk, C. (1996). Effect of MOS on different cecal parameters and on cecal concentrations of enteric bacteria in challenged broiler chicks. *Poultry Sci. Abstract* (in press).
- Stack, R.J. and Cotta, M.A. (1986). Effect of 3-phenylpropanoic acid on growth and cellulose utilization by cellulolytic ruminal bacteria. *Appl. Environ. Microbiol.*, 52: 209-210.
- Stewart, C.S. and Bryant, M.P. (1988). The rumen bacteria. In: *The rumen microbial ecosystem*, Hobson, P.N. (ed.). Elsevier, pp. 21-79.
- Stoll, B.J., Holmgren, J., Bardhan, P.K., Huq, I., Greenhough, W.B., Fredman, P. and Svennerholm, L. (1980). Binding of intraluminal toxin in cholera: trial of GM1 ganglioside charcoal. *Lancet*, ii: 888-891.
- Streeter, M.N., Nisbet, D.J., Martin, S.A. and Williams, S.E. (1994). Effect of malate on ruminal metabolism and performance of steers fed a high concentrate diet. *J. Anim. Sci.*, 72(Suppl. 1): 384.
- Sutton, A.L., Mathew, A.G., Scheidt, A.B., Patterson, J.A. and Kelly, D.T. (1991). Effects of carbohydrate sources and organic acids on intestinal microflora and performances of the weanling pigs. In: *Proc. 6th Int. Symp. on the Digestive Physiology in pigs*, PUDOC, Wageningen, The Netherlands, pp. 422-427.
- Sweet, L.R., Kornegay, E.T. and Lindemann, M.D. (1990). The effect of dietary Luprosil NC on the growth performance and scouring index of weanling pigs. *Agribiol. Res.*, 43: 271-282.
- Takahashi, Y., Kadowaki, K., Tashiro, Y., Takizawa, T. and Kinoshito, T. (1996). Application of fructooligosaccharide to a hemodialysis patient: focused on the change of intestinal bacterial flora. *Bifidus*, 9: 141-150.
- Thacker, P.A., Campbell, G.L. and GrootWassink, J. (1992). The effect of organic acids and enzyme supplementation on the performance of pigs fed barley-based diets. *Can. J. Anim. Sci.*, 72: 395-402.
- Unno, T., Sugawara, M., Nakakuki, T. and Okada, G. (1993). Effect of β -glucan oligosaccharides on the human intestinal microflora. *J. Starch Related Carbohydr. Enz.*, 40: 21-27.
- Yamani, K.A., Ibrahim, H., Rashwan, A.A. and El-Gendy, K.M. (1992). Effects of a pelleted diet supplemented with probiotic (Lacto-Sacc) and water supplemented with a combination of probiotic and acidifier (Acid-Pak 4Way) on digestibility, growth carcass and physiological aspects of weanling New Zealand White rabbits. *J. Appl. Rabbit Res.*, 15:B, 1087-1100.

- Yamazaki, H. and Matsumoto, K. (1994). Purification of *Jerusalem artichoke* fructans and their utilization by bifidobacteria. *J. Sci. Food Agr.*, 64: 461-465.
- Valette, P., Pelenc, V., Djouzi, Z., Andrieux, C., Paul, F., Monsan, P. and Szylił, O. (1993). Bioavailability of new synthesized glucooligosaccharides in the intestinal tract of gnotobiotic rats. *J. Sci. Food Agr.*, 62: 121-127.
- Vanbelle, M., Teller, E. and Focant, M. (1990). Probiotics in animal nutrition: a review. *Arch. Anim. Nutr.*, 40: 543-567.
- Villalba, J.J. and Provenza, F.D. (1996). Preference for flavoured wheat straw by lambs conditioned with intraruminal administrations of sodium propionate. *J. Anim. Sci.*, 74: 2362-2368.
- Voros, C. and Gaal, C. (1992). Effect of *Bacillus* CIP 5832 and enrofloxacin on performance and anaerobic fecal flora of growing rabbits. Deviation of *in vitro* situation from *in vivo* one. *J. Appl. Rabbit Res.*, 15:B, 1153-1159.
- Webb, P.R., Kellogg, D.W., McGahee, M.W. and Johnson, Z.B. (1996). Addition of fructooligosaccharide (FOS) and sodium diacetate (SD) plus decoquinate (D) to milk replacer and starter grain fed to Holstein calves. *J. Dairy Sci.*, 75(Suppl. 1): 300 (Abstr.).
- Wierup, M., Wahlstrom, H. and Engstrom, B. (1992). Experience of a 10-year use of competitive exclusion treatment as part of the *Salmonella* control programme in Sweden. *Int. J. Food Microbiol.*, 15: 287-291.
- Ziprin, R.L. and De Loach, J.R. (1993). Comparison of probiotics maintained by *in vivo* passage through laying hens and broilers. *Poultry Sci.*, 72: 628-635.
- Zoccarato, I., Barbera, S. and Tartari, E. (1995). Effetto dell'impiego di mangime contenente un'associazione antibiotico-probiotico sulle performances del coniglio all'ingrasso. *Zoot. Nutr. Anim.*, 21: 297-304.