

Ecophysiology of annual legumes

Pozo A. del, Aronson J.

in

Sulas L. (ed.).

Legumes for Mediterranean forage crops, pastures and alternative uses

Zaragoza : CIHEAM

Cahiers Options Méditerranéennes; n. 45

2000

pages 223-230

Article available on line / Article disponible en ligne à l'adresse :

<http://om.ciheam.org/article.php?IDPDF=600201>

To cite this article / Pour citer cet article

Pozo A. del, Aronson J. **Ecophysiology of annual legumes**. In : Sulas L. (ed.). *Legumes for Mediterranean forage crops, pastures and alternative uses* . Zaragoza : CIHEAM, 2000. p. 223-230 (Cahiers Options Méditerranéennes; n. 45)



<http://www.ciheam.org/>
<http://om.ciheam.org/>

Ecophysiology of annual legumes

A. Del Pozo¹ and J. Aronson²

¹Facultad de Agronomía, Universidad de Concepción, Casilla 537, Chillán, Chile

²CNRS (UPR 9056), CEFE, 1919, Route de Mende, 34293 Montpellier cedex 5, France

Summary - We review and briefly analyse some of the main ecophysiological attributes found in annual legumes in evolutionary and ecological response to the environmental factors affecting their development, reproduction and survival in Mediterranean Type Ecosystems. We also give some examples of inter- and intra-specific differences that need to be taken into account when introducing or disseminating annual legumes for pasture improvement, rehabilitation efforts, erosion control and/or long term soil restoration programmes.

Key-words: *Medicago*, *Trifolium*, *Ornithopus*, *Biserrula*, reproductive phenology, hardseededness, relative growth rate

Resumé - Les principaux attributs écophysologiques des légumineuses annuelles sont analysés en tant que réponse évolutive et écologique aux facteurs environnementaux affectant leur développement, leur reproduction et leur survie dans les écosystèmes méditerranéens. Quelques exemples de différences inter- et intra-spécifiques sont proposés pour être pris en compte lors de l'introduction ou de la dissémination de légumineuses annuelles pour l'amélioration pastorale, la réhabilitation des parcours, le contrôle de l'érosion ou la restauration des sols.

Mots-clés: *Medicago*, *Trifolium*, *Ornithopus*, *Biserrula*, phénologie de la reproduction, dureté des graines, taux de croissance relatif

Introduction

In mediterranean-type ecosystems (MTEs), vascular plants are exposed to various stresses as the result of interactions with abiotic factors like temperature (especially low and high extremes), excess or insufficient soil water availability (i.e., temporary water logging and prolonged drought), soil acidity or salinity, and nutrient deficiencies. Biotic factors also play a major role in plant ecology and evolution in these regions, including plant-plant interactions (e.g., intra and interspecific competition, neighbor relationships, shading), herbivore impact (e.g., insect defoliation, overgrazing) and mutualist or other kinds of interactions with fungi, bacteria and other micro-organisms. Like all other plants and animals in the MTEs, annual plants must also cope with highly seasonal environments, and a remarkable degree of climatic unpredictability. Under these conditions, being an annual clearly has many advantages, especially in the Mediterranean Basin itself, where annuals may be considered something of a regional 'speciality', often constituting half the regional vegetation whereas they rarely amount to more than one-tenth in other parts of the world (Raven, 1973; Blondel and Aronson, 1999). Clearly, the diversity and abundance of annuals in the Mediterranean region is related to the long history of anthropic disturbances, as pointed out by many authors. Accordingly, it is not surprising how many ruderal and segetal Mediterranean annuals – especially from the eastern part of the Basin and adjacent steppe regions of the Middle East – have successfully colonized California, Chile and, to a lesser extent, south Australia and the Cape Province of South Africa. Among these Mediterranean annual 'invaders' only one group is normally greeted with any enthusiasm: the annual, nitrogen-fixing legumes of high pastoral value such as *Medicago*, *Trifolium* and many others. In addition to various suites of

adaptations allowing survival during periods of prolonged drought, for example, these annual legumes also have remarkable capacity to ‘capitalize’ rapidly on short, intense periods of resource availability and thereby increase their number of offspring dramatically under such conditions.

Despite their numbers, annual Mediterranean legume species can conveniently be considered as a single “functional group” in fields or pastures, i.e., a group of species with similar responses to a given factor, and having a set of common biological (and physiological) attributes that correlate with their overall performance and fitness (Lavorel *et al.*, 1998). However, they do of course differ in many ways, including their responses to a given disturbance or stress, among genera, species and even accessions of a given species. In this short review we briefly analyse the main ecophysiological attributes of annual legumes and some of the environmental factors that affect their development, reproduction and resource allocation patterns, with special emphasis on conditions found in MTEs. We will also give some examples of inter-specific differences that need to be taken into account when introducing or disseminating annual legumes for pasture improvement, rehabilitation efforts, erosion control and/or long term soil restoration programmes.

Plant development and adaptation to seasonality and unpredictability

In the Mediterranean region, and in MTEs in general, annual legumes tend to grow and reproduce during the cool and wet seasons only. In addition to rainfall unpredictability the length of the growing is also very variable in MTEs, and annual legumes show ecotypic differentiation for flowering time in more or less close correlation to the aridity of the collection site (e.g. Piano *et al.*, 1996; Ehrman and Cocks, 1996; Del Pozo *et al.*, 1999a), since increasing aridity and unpredictability go together (Le Houérou, 1984). It has frequently been demonstrated that flowering time plasticity is a common feature of adaptive significance for annuals, including legumes, in arid or semiarid environments including MTEs (Fox, 1989, 1990; Ehrman and Cocks, 1996).

Also the occurrence of summer and early autumn rains in MTEs, which result in false breaks of season, i.e. an early germination of seeds which then die in the absence of follow-up rains, is another environmental constrain for annual plants. To cope this annual legumes show high levels of seed dormancy and mechanism that regulate the pattern of seed germination over more than one growing season. Seed survival in the seed bank is also of great importance to populations; predation by ants and other animals, fungal attack and premature germination of seeds leading to ‘infant mortality’ all impinge on seed bank longevity, and need to be studied carefully for individual species in a given environment (Quinlivan, 1966).

Therefore, the adaptation and long-term persistence of annual legumes in fluctuating environments of MTEs depends largely on the capacity of plants to complete their reproductive cycle, in order to ensure seed production, and on the survival of seeds in the soil over several seasons in what is commonly called a seed-bank.

Seed dormancy

Despite the fact that various mechanisms and components of seed dormancy have been described in legumes, hardseededness (seed coat impermeability) and high-temperature dormancy (*i.e.* the failure of seed to germinate at high temperature when moisture is adequate) are probably the most important ones in Mediterranean legumes. Hardseededness, or impermeability of the seed coat to water, prevents imbibition. This develops during seed maturation and is under both genetic and environmental control. Impermeability is broken by

the influence of diurnal fluctuating temperatures and other natural or artificial processes that scarify the seed coat to produce 'soft' seed (Jansen and Ison, 1995).

Newly ripened seeds of annual legumes have high levels of hardseededness, but the rate of seed softening under field conditions differs greatly among accessions and between species (Table 1). For example, seed softening occurs very rapidly in *Trifolium subterraneum* and *T. isthmocarpum*, but the opposite is true in *Ornithopus compressus* and *Biserrula pelecinus*. In annual medics (*Medicago* spp.) and *T. subterraneum*, the rate of seed softening decreases as depth of burial increases (Taylor and Ewing, 1996); in *O. compressus* and *B. pelecinus* softening occurs more rapidly at 2 cm depth than at 6 cm (Revell *et al.*, 1998; Loi *et al.*, 1999). Buried seeds of annual medics and subterranean clover can survive as hard seeds for up to 12 years without significant loss of viability, increasing in this way seed longevity and legume population persistence (Taylor and Ewing, 1996).

Table 1. Summary of studies reporting weight per seed and seed softening of annual legumes under field conditions.

Species	N° of accessions	Weight per seed (mg)	Hardseededness (%)			Source
			Initial	After 1 summer	After 2 summers	
<i>Medicago</i>						
<i>Littoralis</i>	1	2.2	98	-	33	(1)
<i>Polymorpha</i>	4	2.3-3.0	98-99	-	17-45	(1)
	1		98	85	50	(5)
	34	2-6.6	81-99	30-93	-	(6)
<i>Orbicularis</i>	1	3.9	95	-	75	(1)
<i>Rugosa</i>	1	7.6	96	-	0.5	(1)
<i>Scutellata</i>	4	15.0-17.7	95-98	-	5-30	(1)
<i>Truncatula</i>	8	2.9-4.5	93-99	-	20-60	(1)
<i>Trifolium</i>						
<i>Angustifolium</i>	1	n.a.	91	9	-	(3)
<i>Argutum</i>	1	n.a.	84	44	-	(3)
<i>Cherleri</i>	2	n.a.	93-96	30-47	-	(3)
<i>Clusii</i>	1	n.a.	98	78	-	(3)
<i>Clypeatum</i>	1	n.a.	87	45	-	(3)
<i>Gladuliferum</i>	2	n.a.	98-99	66-73	-	(3)
<i>Lappaceum</i>	1	n.a.	95	70	-	(3)
<i>Isthmocarpum</i>	1	n.a.	90	0.6	-	(3)
<i>Nigrescens</i>	1	n.a.	93	56	-	(3)
<i>Obscurum</i>	1	n.a.	73	32	-	(3)
<i>Purpureum</i>	2	n.a.	85-86	13-23	-	(3)
<i>Subterraneum</i>	1	9.8	98	-	10	(2)
Others						
<i>Biserrula pelecinus</i>	4	0.8-1.6	99.0	-	82-97	(2)
<i>Ornithopus compressus</i>	1	3.5	98.0	-	84-92	(2)
	4	1.7-3.0	96-98	-	35-50	(4)

(1): Lloyd *et al.* (1997); (2): Loi *et al.* (1999); (3): Norman *et al.* (1998); (4): Revell *et al.* (1998); (5): Avendaño *et al.* (1993); (6) Taylor (1996); n.a.: not available

High-temperature dormancy has been detected in various species of *Trifolium* (Jansen and Ison, 1995; Norman *et al.*, 1998). This is an enforced dormancy that occurs when seeds are exposed to temperatures of 30-40 °C. In this range of high temperatures, seeds imbibe water but fail to germinate, even though the same seeds germinate when subsequently placed at 15 °C (Jansen and Ison, 1994). It has been suggested that this type of seed dormancy could play a role in preventing seed from germinating in the case of false break (Jansen and Ison, 1994).

Seed germination

Once seed dormancy has been broken down and soil has enough moisture, the percentage and rate of seed germination depends largely on temperature. The relationship between seed germination and temperature has been studied for various annual legume species, including edible legumes (e.g. Covell *et al.*, 1986; Durmur *et al.*, 1990) and forage legumes (e.g. Jansen and Ison, 1994; Norman *et al.*, 1998). A simple mathematical model has been developed to analyse the germination-temperature relationship and to determine critical temperatures for various species. In this model the rate of seed germination ($1/t$, days⁻¹)- defined as the inverse of the time required to reach a certain percentage of germination -, increases linearly with temperature, from a base temperature (T_b) at which the rate is zero, up to an optimum temperature (T_o), at which this rate is maximal (García-Huidobro *et al.*, 1982; Covell *et al.*, 1986; Del Pozo *et al.*, 1987). The relationship can be described as follows:

$$1/t = a + b T \quad T_b < T < T_o \quad (1)$$

In this model $-a/b$ corresponds to the base temperature and $1/b$ is the thermal time (θ_1 , °C day). However, above the optimum temperature, further increase in temperature reduces $1/t$ until a maximum temperature (T_m) is reached where the rate is again zero (García-Huidobro *et al.*, 1982; Durmur *et al.*, 1990).

Cardinal temperatures (T_b , T_o and T_m) and thermal times for germination of various leguminous species and ecotypes have been determined by fitting linear regressions of rate of germination and sub- and above-optimal temperatures. For Mediterranean legumes, base temperatures ranged from 0 to 5 °C, optimum temperature ranged from 16 to 22 °C, whereas maximum temperature ranged from 30 to 40 °C (Covell *et al.*, 1986 for edible legumes; Norman *et al.*, 1998 for forage legumes).

Flowering time

As mentioned above, annual legumes show both ecotypic differentiation and a high degree of plasticity in flowering time. Although severe water deficits may affect the development of annual legumes, time to flowering is mainly influenced by genotype, temperature and photoperiod (Roberts *et al.*, 1997). Plants from MTEs are usually long-day plants, i.e. flowering is delayed when daylengths are shorter than a critical photoperiod (P_c).

The influence of temperature and photoperiod on flowering time has been studied in various leguminous taxa including *Medicago* spp. (Clarkson and Russell, 1975; 1979; Del Pozo *et al.*, 1999a) and *Trifolium* spp. (Evans *et al.*, 1992).

The analysis has been made using a linear model which relate the rate of progress to flowering ($1/f$) - defined as the inverse of time from emergence to flowering (f) - with mean diurnal temperature (T , °C) and mean photoperiod (P , h day⁻¹) (Summerfield *et al.*, 1991; Evans *et al.*, 1992; Del Pozo *et al.*, 1999a), according to the equation:

$$1/f = a' + b'T + c'P \quad (2)$$

where a' , b' and c' are constants specific to each accession (Summerfield *et al.*, 1991).

Studies of flowering responses to temperature and photoperiod conducted on eight cultivars of *Trifolium subterraneum* grown in Australia (Evans *et al.*, 1992), and nine accessions of *M. polymorpha* collected from a desert-mediterranean climatic gradient in Chile (Del Pozo *et al.*, 1999a), demonstrated a great variability in flowering time. In addition, in both studies the parameters b' and c' of model 2 differed among accessions, clearly indicating the existence of genotypic variation for sensitivity to both temperature and photoperiod. Detailed genetic studies are now underway to elucidate these differences and render them amenable to utilisation by breeders.

Growth and allocation

Legumes usually have higher nitrogen concentration in both vegetative and reproductive organs than grasses or other dicotyledenous plants (e.g. Del Pozo, 1992; McKey, 1994; Navas *et al.* 1995). According to McKey (1994), nitrogen-rich leaves of legumes should allow higher rates of photosynthesis, especially at high light intensities, and therefore favor rapid growth of leaves and plant production, during favourable periods.

However, several studies have shown that the high N content in vegetative organs of legumes does not lead to a higher relative growth rate (RGR) as compared with annual grasses and other families. For example, Elias and Chadwick (1979) compared the early seedling growth of 28 grasses and 12 legumes (all cultivated pasture species), and found that the relative growth rate of the legumes was slightly lower than that of the grasses (0.156 d^{-1} vs. 0.171 d^{-1}). Similarly, in a study comparing the growth of 27 annual species of three botanical families occurring in Mediterranean grasslands, Marañón and Grubb (1993) reported that the eight legume taxa tested showed lower relative growth rate than the 12 grasses or the seven Asteraceae studied.

In contrast, a literature review for over 20 crop species indicated that the ratio between standing dry mass and nitrogen content (*i.e.*, nitrogen use efficiency, NUE), is lower in legumes than in C_3 grasses, suggesting that important differences in physiological characteristics may exist between the two groups, including the nitrogen and carbon allocation patterns, growth and rates of photosynthesis under varying conditions, between legumes and grasses (Del Pozo *et al.*, 1999b).

During the early stages of growth, relative growth rate (RGR) has often been found to be proportional to mean plant organic-N concentration (PNC: Ågren, 1985; Greenwood *et al.* 1991):

$$RGR = NP \times PNC \quad (3)$$

where NP is plant nitrogen productivity ($\text{gDW mol N}^{-1} \text{ d}^{-1}$). NP thus expresses the way internal nitrogen is used for growth. More recently Garnier and Vancaeyzeele (1994) have shown that NP can be factored into two terms:

$$NP = LNR \times N_LP \quad (4)$$

where LNR is leaf nitrogen ratio (ratio between leaf and total plant nitrogen contents) and N_LP is the leaf nitrogen productivity, *i.e.* ratio of growth rate to leaf nitrogen concentration ($\text{gDW mol N}^{-1} \text{ d}^{-1}$).

Although there are few direct comparisons between legumes and non-legumes for these traits, it does seem that legumes have a lower nitrogen productivity and a greater x-intercept (i.e., the minimum N concentration where RGR is equal to zero) than non-legumes such as C₃ cereal grasses (Table 2). Furthermore, differences in nitrogen productivity between the two groups of species were mainly explained by differences in leaf nitrogen productivity (Table 2). This latter trait has been shown to be related to photosynthetic nitrogen use efficiency (PNUE), i.e. the ratio between rate of photosynthesis and leaf nitrogen concentration (Garnier *et al.*, 1995). In fact, PNUE was found to be lower in legumes than in cereals (Del Pozo *et al.*, 1999b). It seems that the high nitrogen concentration in legumes may have more to do with a high nitrogen demand for seed production at a time when nitrogen fixation is shut off than with a high production potential (Del Pozo *et al.*, 1999b).

Table 2. Plant nitrogen concentration (PNC), relative growth rate (RGR), nitrogen productivity (NP) and its components in annual C₃ grasses and legume plants. LNR and N_LP are the leaf nitrogen ratio and leaf nitrogen productivity, respectively (see text for definition of parameters). Source: Del Pozo *et al.* (1999b).

Species	PNC (mmol N g ⁻¹)	RGR (g g ⁻¹ d ⁻¹)	NP (g mol ⁻¹ d ⁻¹)	LNR (mol mol ⁻¹)	N _L P (g mol ⁻¹ d ⁻¹)
<i>Medicago minima</i>	2.70	0.055	19.8	0.66	30
<i>Vicia faba</i>	3.00	0.031	10.3	0.38	27.2
<i>Bromus madritensis</i>	1.94	0.061	30.6	0.51	60
<i>Triticum aestivum</i>	2.61	0.070	26.8	0.31	85.3

Concluding remarks

In our introduction, we suggested that despite their numbers, annual Mediterranean legumes could be considered as a single “functional group”. Indeed, when compared to annual C₃ grasses, or other large but coherent groups of annuals, e.g., Asteraceae, this assertion appears to hold up. More detailed studies of specific responses to specific stresses are required to test this idea further, both in monocultures and, especially, in mixed cultures. However, we have also given examples of inter- and intra-specific differences that can be useful when introducing or disseminating annual legumes for pasture improvement, rehabilitation efforts, erosion control and/or long term soil restoration programmes. For example, flowering responses to temperature and photoperiod in *Trifolium subterraneum* and *Medicago polymorpha* accessions showed great variability in flowering time, which is a variable of critical importance to pasture management in seasonal, highly unpredictable environments such as most MTEs. Similar variability is found in winter vigour, seed dormancy components and other agronomically important traits. More research is needed on genetic components of intra- and inter-specific variability in response to environmental conditions and, of course, performance of selected ecotypes or genotypes under field conditions.

In practical terms, one promising strategy for range managers is to develop appropriate mixtures of species and accessions for combined use. In this way, as Piano *et al.* (1996), Del Pozo *et al.* (1999b) and others have emphasized, it should be possible to increase chances of high productivity and at least some production of seeds regardless of the rainfall and temperature patterns of the current year.

References

- Ågren, G.I. (1985). Theory for growth of plants derived from the nitrogen productivity concept. *Physiol. Plantarum*, 64: 17-28.
- Avendaño, J., Del Pozo, A., Ovalle, C. (1993). Hardseededness under field condition of *Medicago polymorpha* in the mediterranean zone of Chile. Proc. 17th Int. Grass. Cong., Hamilton, New Zealand, pp. 2206-2207.
- Blondel, J., Aronson, J. (1999). *Biology and wildlife in the Mediterranean region*. Oxford University Press, Oxford, UK.
- Clarkson, N.M., Russell, J.S. (1975). Flowering responses to vernalization and photoperiod in annual medics (*Medicago* spp.). *Aust. J. Agric. Res.*, 26 : 831-838.
- Clarkson, N.M. and Russell, J.S. (1979). Effect of temperature on the development of two annual medics. *Aust. J. Agric. Res.*, 30: 909-916.
- Covell, S., Ellis, R., Roberts, E.H. and Summerfield, R.J. (1986). The influence of temperature on seed germination rate in grain legumes. *J. Exp. Bot.*, 37:705-715.
- Del Pozo, A. (1992). *Carbon/Nitrogen Relations in Cereals and Legumes*. Ph.D. Thesis, University of Reading, United Kingdom.
- Del Pozo, A., García-Huidobro, J.; Novoa, R., Villaseca, S. (1987). Relationship of base temperature to development of spring wheat. *Exp. Agric.*, 23: 21-30.
- Del Pozo, A., Ovalle, C., Aronson, J., Avendaño, J. (1999a) Developmental responses to temperature and photoperiod in ecotypes of *Medicago polymorpha* L. along an environmental gradient in central Chile. *Ann. Bot.* (in press).
- Del Pozo, A., Garnier, E., Aronson, J. (1999b) Contrasted nitrogen utilization in annual C₃ grasses and legume crops: physiological explorations and ecological consequences. *Acta Oecologica* (in press).
- Dumur, D., Pilbeam, C., Craigon, J. (1990). Use of Weibull function to calculate cardinal temperatures in faba bean. *J. Exp. Bot.*, 41: 1423 -1430.
- Ehrman, T., Cocks, P.S. (1996). Reproductive patterns in annual legume species on an aridity gradient. *Vegetatio*, 122:47-59.
- Elias, C.O., Chadwick M.J. (1979). Growth characteristics of grass and legume cultivars and their potential for land reclamation. *J. Appl. Ecol.*, 16:537-544.
- Evans, P.M., Lawn, R.J., Watkinson, A.R. (1992). Use of linear models to predict the date of flowering in cultivars of subterranean clover. *Aust. J. Agric. Res.*, 43:1547-1558.
- Fox, G.A. (1989). Consequences of flowering-time variation in a desert annual: Adaptation and history. *Ecology*, 70:1294-1306.
- Fox, G.A. (1990). Drought and the evolution of flowering time in desert annuals. *Amer. J. Bot.*, 77: 1508-1518.
- García-Huidobro, J., Monteith, J.L., Squire, G.R. (1982). Temperature and germination of pearl millet (*Pennisetum typhoides* S.& H.) I. Constant temperature. *J. Exp. Bot.*, 33: 288-296.
- Garnier, E., Vancaeyzeele, S. (1994). Carbon and nitrogen content of congeneric annual and perennial grass species: relationships with growth. *Plant, Cell Environ.*, 17:399-407.
- Garnier, E., Gobin, O., Poorter, H. (1995). Nitrogen productivity depends on photosynthetic nitrogen use efficiency and on nitrogen allocation within the plant. *Ann. Bot.*, 76:667-672.
- Greenwood, D.J., Gastal, F., Lemaire, G., Draycott, A., Millard, P., Neeteson, J.J. (1991). Growth rate and %N of field grown crops: theory and experiments. *Ann. Bot.*, 67:181-190.
- Jansen, P.I., Ison, R.L. (1994). Temperature effects on germination of *Trifolium balansae* and *Trifolium resupinatum* with special reference to high-temperature dormancy. *Aust. J. Agric. Res.*, 45: 689-701.
- Jansen, P.I., Ison, R.L. (1995). Factors contributing to the loss of seed from the seed-bank of *Trifolium balansae* and *Trifolium resupinatum* over summer. *Aust. J. Ecol.*, 20: 248-256.

- Lavorel, S., Touzard, B., Lebreton, F-D, Clément, B. (1998). Identifying functional groups for response to disturbance in an abandoned pasture. *Acta Oecologia*, 19: 227-240.
- Le Houérou, H.N. (1984). Rain use efficiency: a unifying concept in arid-land ecology. *J. Arid Environ.*, 7: 213-247.
- Lloyd, D.L., Taylor, G.B., Johnson, B., Teasdale, K.C. (1997). Patterns of seed softening and seedling emergence of nineteen annual medics during three years after a single seed crop in southern Queensland. *Aust. J. Exp. Agric.*, 37: 767-778.
- Loi, A., Cocks, P.S., Howieson, J.G., Carr, S.J. (1999). Hardseededness and the pattern of softening in *Beserrula pelecinus* L., *Ornithopus compressus* L., and *Trifolium subterraneum* L. seeds. *Aust. J. Agric. Res.*, 50: 1073-1081.
- McKey, D. (1994). Legumes and nitrogen: the evolutionary ecology of a nitrogen-demanding lifestyle. in: Sprent, J.J., McKey, D: (Eds.), *Advances in Legume Systematics, Vol. 5: The Nitrogen Factor*, Royal Botanic Gardens, Kew. pp. 211-228.
- Marañón, T., Grubb, P.J. (1993). Physiological basis and ecological significance of the seed size and relative growth rate relationship in Mediterranean annuals. *Funct. Ecol.*, 7: 591-599.
- Navas, M.L., Guillerm, J.L., Fabreguettes J., Roy J. (1995). The influence of elevated CO₂ on community structure, biomass and carbon balance of mediterranean old-field microcosms. *Global Change Biol.*, 1: 325-335.
- Norman, H.C., Cocks, P.S., Smith, F.P., Nutt, B.J. (1998). Reproductive strategies in Mediterranean annual clovers: germination and hardseededness. *Aust. J. Agric. Res.*, 49: 973-982.
- Piano, E., Pecetti, L., Carroni, A. (1996). Climatic adaptation in subterranean clover populations. *Euphytica*, 92:39-44.
- Quinlivan, B.J. (1966). The relationship between temperature fluctuations and the softening of hard seeds of some legume species. *Aust. J. Agric. Res.*, 36: 119-144.
- Raven, P.H. (1973). The evolution of Mediterranean floras. In: *Mediterranean-Type Ecosystems, Origin and Structure*. di Castri, F. and Mooney, H.A. (eds.). Springer, Heidelberg. 213-224.
- Revell, C.K., Taylor, G.B., Cocks, P.S. (1998). Long-term softening of surface and buried hard seeds of yellow serradella grown in a range of environments. *Aust. J. Agric. Res.*, 49: 673-685.
- Roberts, E.H., Summerfield, R.J., Ellis, E.H., Craufurd, P.Q., Wheeler, T.R. (1997). The induction of flowering. In: *The Physiology of Vegetable Crops*. H.C. Wien (ed.). CAB International. 69-99.
- Summerfield, R.J., Roberts, E.H., Ellis, R.H., Lawn, R.J. (1991). Toward the reliable prediction of time to flowering in six annual crops. I. The development of simple models for fluctuating field environments. *Exp. Agric.*, 27:11-31.
- Taylor, G.B., Ewing, M.A. (1996). Effects of extended (4-12 years) burial on seed softening in subterranean clover and annual medics. *Aust. J. Exp. Agric.*, 36: 145-150.
- Taylor, G.B. (1996). Incidence and measurement of autumn seed softening within *Medicago polymorpha* L. *Aust. J. Agric. Res.*, 47: 575-586.